

Perturbation technique to retrieve scattering medium stratification

Igor N. Polonsky

School of Physics,
University of New South Wales
Sydney, 2052, Australia

E-mail: polonsky@phys.unsw.edu.au

Michael A. Box

School of Physics,
University of New South Wales
Sydney, 2052, Australia

E-mail: M.Box@unsw.edu.au

Abstract

It is well known that optical satellite remote sensing is mostly based on measurements of the radiance exiting the top of the Earth's atmosphere and is interpreted using a comparison of the data with pre-calculated tables for some "pure" aerosol models. However, such an approach seems to meet some difficulties connected to both the necessity to search in the multi-dimensional parameter space and increasing the volume of the pre-calculated database, if the number of basic aerosol components increases. This problem becomes more real now because modern satellite sensors (MISR, POLDER) provide more information by performing multi angle radiance measurements over the same pixel and more detailed characteristics of the atmospheric aerosol profile (not only the aerosol optical depth and Angstrom coefficient) are expected to be retrieved.

Recently, it have been shown that the perturbation technique was an ideal tool to solve such atmospheric physics problems as an investigation of the effect of a variation in aerosol profile on fluxes. Moreover, it has shown its efficiency in analysis of the optical remote sensing in the simplest case of sounding of a homogeneous medium.

In the present paper the perturbation technique is applied to a realistic atmosphere-ocean model. We performed our simulations for a stratified slab medium with an underlying surface. Perturbations considered include variations of the profiles of extinction coefficient and single scattering albedo, and also the adding of a small cloud contamination. It was shown that the perturbation technique allows predicting the effect of such the variations of the atmospheric profile to the exiting radiance with high accuracy. The nature of the errors is analyzed and discussed.

Perturbation technique to retrieve scattering medium stratification

Igor N. Polonsky, Michael A. Box

1. Introduction

It is well known that the atmospheric aerosol exerts a significant impact on the Earth's climate, both directly (d'Almeida et al., 1991), by absorbing and scattering radiation, and indirectly (Charlson et al., 1992; Curry, 1995) by their effect on cloud optical and microphysical properties. To better understand these effects it is necessary to know not only the aerosol optical depth, but also the aerosol scattering and absorption properties, and their vertical profiles. In other words, an adequate model of global aerosol distribution, taking into account their seasonal and spatial variability, has to be developed.

The integrated field experiments being conducted (like ASTEX/MAGE in June 1992 (Huebert et al., 1996), SCAR-B in August-September 1995 (Kaufman et al., 1998), ACE-1 in November-December 1995 (Bates et al., 1988), TARFOX in July 1996 (Russell, 1999)) can specify many features of local aerosol atmosphere models and their results are the best touchstone for any kind of retrieval algorithms, especially for satellite remote sensing.

The only approach that could provide the necessary spatial and temporal coverage is Earth observation from space. During the last five years several sensors to fulfill this task have been launched. Moreover, some of these modern sensors provide much more information than those of previous generations, by viewing the same surface area under different angles (POLDER on board ADEOS (1996) has up to 14 angles (Deschamps et al., 1994), MISR on EOS (1999) has 9 angles (Diner et al., 1991)). Substantial increasing of the available information makes it possible to retrieve not only the aerosol optical depth, but also aerosol stratification, detection of absorbing aerosol (over the oceans) or estimation of the bi-directional reflection function of the surface (over the land) (Kaufman et al., 1997). However, this leads to the retrieval algorithms becoming more sophisticated and causes the number of base aerosol models to increase.

Nevertheless, the actual number of aerosol models which are really taken into account is usually limited to constrain the size of the pre-calculated database (Martonchik et al., 1998). This could be a possible source of retrieval error, if non-standard aerosol contributions occur, or in the case of an undetected small cloud contamination of a pixel, but no thorough sensitivity study has yet been performed because of the multiple factors nature of the problem. However, the papers by Mishchenko and Travis (Mishchenko and Travis, 1997a, 1997b) should be noted in which, despite of the simplest aerosol-ocean model being considered, the important conclusion that the algorithm based on multiple viewing angle measurements performs far better than that based on single viewing angle radiance measurements is made.

Usually, to simulate satellite remote sensing the approach based on developing of pre-calculated database (Rao et al., 1989; Mishchenko and Travis, 1997a; Martonchik et al., 1998) is used. In this paper a fundamentally different technique is applied to the problem of multiangle satellite observation of a stratified atmosphere. This approach, known as the perturbation technique, has been adapted from nuclear reactor theory (Marchuk, 1964; Gerstl, 1980), and developed comprehensively for the purpose of the radiative transfer in atmosphere in the papers of Box and his co-authors (Box et al., 1988a, 1988b, 1989a, 1989b; Trautmann et al., 1992). This approach shows its efficiency to analyze the possibility of the retrieval of the optical thickness, single scattering albedo and phase function (Sendra and Box, 2000) in the case of the sounding of a homogeneous medium. In this paper it will be applied to the sounding of a stratified medium. The perturbation approach also seems to be a perfect base to develop the retrieval algorithm, because it provides the necessary derivatives and hence the standard techniques of the optimization of multivariable non-linear functions (Fletcher, 1987) may be successfully implemented.

2. Perturbation technique

The perturbation technique of radiative transfer theory is based on the joint solution of both the direct and adjoint equations, and has been used for more than fifty years. Initially it was successfully applied to nuclear reactor problems (Lewins, 1965; Bell and Glasstone, 1970). Marchuk (Marchuk, 1964) was the first who has introduced this approach as a general technique to interpret optical measurements into the field of the atmospheric optics. However, the most successful applications of the perturbation technique started more than a decade later. This approach was applied to compute atmospheric radiative effects (Gerstl, 1980; Box et al., 1988a, 1989a, 1989b; Trautmann et al., 1992), to consider the problems of thermal sounding (Ustinov, 1990) and the photometric observation of solar radiation reflected from the optically thick vertically inhomogeneous planet atmosphere (Ustinov, 1991a, 1991b, 1992), and to investigate the sensitivity of exiting radiances to optical characteristics in the simplest case of a homogeneous atmosphere (Box and Sendra, 1995).

Let us introduce briefly the basic results. We consider only the case of radiation transfer in a plane parallel horizontally homogeneous atmosphere with solar illumination. In the most common cases, the radiative effects (for example, fluxes) or optical measurements are given by

$$E = \iint I(z, \Omega) R(z, \Omega) dz d\Omega, \quad (1)$$

where $I(z, \Omega)$ is the radiance at altitude z and in the direction Ω , $R(z, \Omega)$ is the response function which to simulate satellite measurements has the form:

$$R(z, \Omega) = \delta(z - z_t) \delta(-\mu - \mu_R) \delta(\phi - \phi_R), \quad (2)$$

where μ is the cosine of the zenith angle, $\delta(z)$ is the Dirac δ -function, z_t is the altitude of the atmosphere “top”, μ_R and ϕ_R are the zenith angle cosine and azimuth angle of the satellite receiver.

$I(z, \Omega)$ satisfies the radiative transfer equation (RTE)

$$\left[\mu \frac{d}{dz} + \varepsilon(z) \right] I(z, \Omega) = \frac{\sigma(z)}{4\pi} \iint \chi(z, \Omega, \Omega') I(z, \Omega') d\Omega' + Q(z, \Omega), \quad (3)$$

where $\varepsilon(z)$ and $\sigma(z)$ are the extinction and scattering coefficients, respectively, $Q(z, \Omega)$ is the source function, which has the form

$$Q(z, \Omega) = S_0 \delta(z - z_t) \delta(\mu - \mu_0) \delta(\phi - \phi_0),$$

here μ_0 and ϕ_0 are the zenith angle cosine and azimuth angle of the Sun. $\chi(\Omega, \Omega')$ is the phase function, normalized by

$$\frac{1}{4\pi} \iint \chi(\Omega, \Omega') d\Omega' = 1.$$

For the sake of simplicity it is convenient to rewrite equation (4) in the operator form

$$\hat{L}I(z, \Omega) = Q(z, \Omega), \quad (4)$$

where

$$\hat{L} = \mu \frac{d}{dz} + \varepsilon(z) - \frac{\sigma(z)}{4\pi} \iint d\Omega' \chi(z, \Omega, \Omega') \otimes. \quad (5)$$

The notation \otimes is used to indicate that the final term is an integral operator, not a simple definite integral.

Let us introduce an adjoint operator \hat{L}^+ , which is defined by requiring that

$$\langle I^+ (\hat{L} I) \rangle = \langle (\hat{L}^+ I^+) I \rangle, \quad (6)$$

where $\langle \dots \rangle = \iint (\dots) dz d\Omega$ denotes a scalar product and I^+ is the solution of the adjoint equation

$$\hat{L}^+ I^+(z, \Omega) = R(z, \Omega). \quad (7)$$

Taking into account the common boundary conditions for the radiance (Lenoble, 1985, p.179),

i.e.

$$\begin{aligned} I(z=0, \Omega) &= \frac{1}{\pi} \int_0^{2\pi} d\phi' \int_{-1}^0 d\mu' \mu' \rho(\Omega, \Omega') I(z=0, \Omega'), & \mu > 0, \\ I(z=z_t, \Omega) &= 0, & \mu < 0, \end{aligned} \quad (8)$$

where $\rho(\Omega, \Omega')$ is the bi-directional reflection function of the underlying surface and the sunlight is taken into account in the radiative transfer equation itself, we can obtain from (6) the explicit form for \hat{L}^+ (Marchuk, 1964; Box et al. 1988a; Ustinov, 1991a)

$$\hat{L} = -\mu \frac{d}{dz} + \varepsilon(z) - \frac{\sigma(z)}{4\pi} \iint d\Omega' \chi(z, \Omega, \Omega') \otimes. \quad (9)$$

and the boundary condition for I^+

$$\begin{aligned} I^+(z=0, \Omega) &= \frac{1}{\pi} \int_0^{2\pi} d\phi' \int_0^1 d\mu' \mu' \rho(\Omega', \Omega) I(z=0, \Omega'), & \mu < 0, \\ I^+(z=z_r, \Omega) &= 0, & \mu > 0. \end{aligned} \quad (10)$$

Let us consider two atmospheres whose optical properties differ slightly (the first one we denote as “base” and the other as “perturbed”) and let

$$\hat{L}_p = \hat{L}_B + \delta\hat{L}, \quad (11)$$

It can be shown that if only the most significant contributions are taken into account the effect perturbation δE which corresponds to the perturbation $\delta\hat{L}$ of the atmosphere is given by (Marchuk, 1965; Box et al., 1988a; Ustinov, 1991a)

$$\delta E = -\langle I^+ \delta\hat{L} I \rangle. \quad (12)$$

More detailed results and discussion concerning higher-order terms can be found in (Box et al., 1988b). Unfortunately, it is difficult to provide simple criteria to determine whether a given atmosphere perturbation is small enough and hence to estimate the accuracy of (12), but some simulation results provided below will serve to give a rough idea.

3. Simulation technique

We introduce a Cartesian coordinate system with the z-axis directed to atmosphere “top” along the normal to its boundary and the x-axis in the plane of sunlight incidence. The direction of the light propagation is specified by $\Omega = (\vartheta, \phi)$, where ϑ is the zenith angle measured from the positive z-axis and ϕ is the azimuth angle measured clockwise from the positive x-axis.

To perform the perturbation calculation we must evaluate integral (12). To simplify the integration, let both $I(z, \Omega)$ and $I^+(z, \Omega)$ be represented as a series of associated Legendre functions:

$$I(z, \Omega) = \sum_{n=0}^{\infty} \sum_{m=0}^n A_{nm} \Psi_{nm}(z) P_n^m(\mu) \cos(m\phi), \quad (13)$$

$$I^+(z, \Omega) = \sum_{n=0}^{\infty} \sum_{m=0}^n A_{nm} \Psi_{nm}^+(z) P_n^m(-\mu) \cos(m\phi).$$

Here, $A_{nm} = (2n+1)(2-\delta_{0m})$, $\mu = \cos(\vartheta)$, and $\chi(z, \beta)$ ($\beta = (\Omega, \Omega')$ is the scattering angle) may also be expanded in a series of Legendre polynomials:

$$\chi(z, \beta) = \sum (2n+1) \chi_n(z) P_n(\beta). \quad (14)$$

Note that in the most common case the atmosphere perturbation can be represented in the form

$$\begin{aligned} \delta \hat{L}(z, \Omega) = & \delta \sigma_e(z) \left[1 - \frac{\omega_0(z)}{4\pi} \int d\Omega' \chi(z, \Omega, \Omega') \otimes \right] \\ & - \sigma_e(z) \frac{\delta \omega_0(z)}{4\pi} \int d\Omega' \chi(z, \Omega, \Omega') \otimes, \\ & - \sigma_e(z) \frac{\omega_0(z)}{4\pi} \int d\Omega' \delta \chi(z, \Omega, \Omega') \otimes \end{aligned} \quad (15)$$

where the first member is due to the extinction coefficient variation $\delta \sigma_e$, the second corresponds to the single scattering albedo variation $\delta \omega_0$ and the third describes the influence of the phase function variation $\delta \chi$. Therefore, the corresponding expression for the effect perturbation can be obtained by substitution of (13-15) into (12), and takes the form

$$\delta E = \sum_{n=0}^{\infty} \sum_{m=0}^n (-1)^n A_{nm} \cos(m\phi_R) \int \Psi_{nm}^+ \Psi_{nm} \{ [1 - \omega_0 \chi_n] \delta \sigma_e - \sigma_e \chi_n \delta \omega_0 - \sigma_e \omega_0 \delta \chi_n \} dz, \quad (15)$$

It is interesting to point out that if we need to estimate the variation of the exiting radiances due to adding some aerosol characterized by the parameters $\sigma_e^p, \sigma_s^p, \chi^p$, which are independent of altitude, Eq.15 is simplified to

$$\delta E = \sum_{n=0}^{\infty} \sum_{m=0}^n (-1)^n A_{nm} \cos(m\phi_R) [1 - \omega_0^p \chi_n^p] \sigma_e^p \int \psi_{nm}^+ \psi_{nm} dz, \quad (16)$$

and the integral over z should be calculated only once, even if the influence of different aerosols is to be investigated.

The natural way to calculate ψ_{nm} is the spherical harmonics approximation (SHA) (Dave, 1975). Moreover, it allows one to get both ψ_{nm} and ψ_{nm}^+ during the same calculation without considerable expense of computer time. It is well known that SHA allows all integrals from the radiance over angles to be estimated with high accuracy, but unavoidable oscillations in the radiance angle distribution appear near the boundary. To improve the accuracy the iteration procedure (Lenoble, 1985, p.35) is used. This idea is based on using the formal solution of the radiative transfer equation (4), which has the form

$$\begin{aligned} I(z, \Omega) &= I_{SHA}(z_T, \Omega) \exp\left[-\frac{\tau(z)}{\mu}\right] + \frac{1}{\mu} \int_z^{z_T} \exp\left[-\frac{\tau(z) - \tau(z')}{\mu}\right] J(z', \Omega) dz', & \mu > 0 \\ I(z, \Omega) &= J(z, \Omega), & \mu = 0 \\ I(z, \Omega) &= I_{SHA}(0, \Omega) \exp\left[-\frac{\tau(z) - \tau(z_i)}{\mu}\right] - \frac{1}{\mu} \int_0^z \exp\left[-\frac{\tau(z) - \tau(z')}{\mu}\right] J(z', \Omega) dz', & \mu < 0 \end{aligned} \quad (17)$$

where the optical thickness $\tau(z)$ is defined in the form

$$\tau(z) = \int_z^{z_T} \sigma_e(z) dz$$

and the SHA solution $I_{SHA}(z, \Omega)$ is being used to get

$$J(z, \Omega) = \frac{\sigma(z)}{4\pi} \iint \chi(z, \Omega, \Omega') I_{SHA}(z, \Omega') d\Omega'. \quad (18)$$

There are two important reasons to improve the SHA solution. Firstly, it is necessary to simulate satellite measurements. Secondly, to estimate some effects of the perturbation the integral

$$\delta E = - \int_0^{z_T} dz \int_{4\pi} d\Omega I^+(z, \Omega) I(z, \Omega) \quad (19)$$

needs to be calculated. If we take into account that both $I(z, \Omega)$ and $I^+(z, \Omega)$ are contributed significantly by the singular component $S^{(+)} \exp[-\tau(z)/\mu_0] \delta(\Omega - \Omega_0)$ at small z (“0” denotes either the Sun illumination or receiver angles), we meet again the necessity of the exact radiance calculation.

4. Numerical results

We consider the widely used model of the atmosphere as a stratified slab with an underlying surface. Although there is no restriction on the reflection properties of the underlying surface, for the sake of simplicity we consider the Lambertian case. Moreover, it is a reasonable approximation to model the real atmosphere-ocean system. In these simulations three profiles (Lenoble, 1985) are chosen as appropriate for our purpose. The characteristics of the layers are given in Table 1. Note that Model III is very similar on Model I except for the large cloud. We introduce this model to check the applicability of the perturbation technique in the case of an optically thick atmosphere. The US standard atmosphere is chosen as a model of the molecular atmosphere. We restrict our simulation to a single wavelength of 0.55 μm , at which the contribution of the molecular atmosphere cannot be neglected, but it is still less than the aerosol component.

Let us consider a typical problem of the aerosol satellite remote sensing. We would like to investigate how the exiting radiances change if the basic characteristics of the lowermost layer, such as the optical thickness or the single scattering albedo, are changed. Also we are interested to estimate the influence of a small cloud contamination to it. Let us solve this problem using two approaches: the perturbation technique and the direct solution of the RTE, which is the common approach to do it. Taking into account that the perturbation technique assumes a linear dependence of the effect on the perturbation (12), it is convenient to evaluate the corresponding derivatives instead of the absolute values. Moreover, they describe the sensitivities of the exiting radiance to variation of these parameters.

Using the perturbation technique we calculate the following characteristics:

$$\begin{aligned}\frac{dI}{d\sigma_e} &= \sum_{n=0}^{\infty} \sum_{m=0}^n (-1)^n A_{nm} \cos(m\phi_R) \int \Psi_{nm}^+ \Psi_{nm} [1 - \omega_0 \chi_n] dz, \\ \frac{dI}{d\omega_0} &= \sum_{n=0}^{\infty} \sum_{m=0}^n (-1)^n A_{nm} \cos(m\phi_R) \int \Psi_{nm}^+ \Psi_{nm} \chi_n dz, \\ \frac{dI}{d\tau_C} &= \sum_{n=0}^{\infty} \sum_{m=0}^n (-1)^n A_{nm} \cos(m\phi_R) \int \Psi_{nm}^+ \Psi_{nm} [1 - \omega_0^C \chi_n^C] dz,\end{aligned}\tag{20}$$

where τ_C is the cloud optical thickness, ω_0^C is the cloud single scattering albedo and χ_n^C are the expansion coefficients of the cloud phase function in Legendre polynomials. The direct solution of the RTE provides us with the corresponding estimates

$$\begin{aligned}\frac{\Delta I}{\Delta\sigma_e} &= \frac{I(z, \Omega; \sigma_e + \Delta\sigma_e) - I(z, \Omega; \sigma_e)}{\Delta\sigma_e}, \\ \frac{\Delta I}{\Delta\omega_0} &= \frac{I(z, \Omega; \omega_0 + \Delta\omega_0) - I(z, \Omega; \omega_0)}{\Delta\omega_0}, \\ \frac{\Delta I}{\Delta\tau_C} &= \frac{I(z, \Omega; \tau_C + \Delta\tau_C) - I(z, \Omega; \tau_C)}{\Delta\tau_C}.\end{aligned}\tag{21}$$

We performed the calculation assuming the sensor azimuth angle is 90° and the surface albedo is 0.05 for several values of $\Delta\sigma_e$, $\Delta\omega_0$ and $\Delta\tau_C$. This allows us to estimate the applicability range of the perturbation technique over the parameters variation area.

The results of our calculation for the atmosphere models are given in Tables 2-9. For convenience they also contain the relative error, which, for example, for the extinction coefficient perturbation has the form

$$\text{relative error} = \frac{\frac{dI}{d\sigma_e}}{\Delta I} - 1\tag{22}$$

Tables 2-4 show how the perturbation technique can predict the effect of the extinction coefficient variation. The range was chosen to be rather wide from $-0.5\sigma_e$ to σ_e (that is, from

half the original value to double it). Tables 5-7 contain results for the single scattering albedo variations. Tables 8-9 show the sensitivity of the exiting radiance to a relatively small cloud contamination.

In all the tables we can see that the linear approximation (12) of the dependence of the effect on the parameter perturbation is reasonable in most of the case considered, especially for atmosphere models I and II, which are not optically thick. The tables give us the area of the perturbation technique applicability. For distinctness let us set 10% as a permissible limit of the relative error. We can see that for atmosphere model I, which describes clear atmosphere, the perturbation technique and direct simulation results coincide very well in practically all considered cases, except for a very few. This can be explained by the small multiple scattering contribution to the exiting radiances. For model II, which corresponds to a rather polluted atmosphere, the accuracy of the perturbation technique estimation becomes unacceptable at extinction coefficient variation $|\Delta\sigma_e| \gg 0.1\sigma_e$, but is still tolerable for all considered variations of the single scattering albedo, except $\Delta\omega_0 = -0.5$.

Consideration of the results for model III shows that we meet an interesting situation, where noting that the optical thickness of the lowermost layer is not small ($\tau=10$), and its single scattering albedo is close to 1 ($\omega_0=0.998$), it is clear that multiple scattering contributes significantly to the exiting radiance, but the nonlinearity of its dependence on the extinction coefficient variation decreases slightly in comparison to model II. In contrast, Table 7 shows us that to predict the effect of the single scattering albedo variation on the exiting radiance we can use the perturbation technique only at $|\Delta\omega_0| < 0.01$ to meet the above criteria. Are there some contradictions? Let us recall some analytical results of the asymptotic theory of an optically thick weakly absorbing medium. For this case the exiting radiance for a semi-infinite layer is given by (Zege, 1991)

$$I(z_t, \Omega; \Omega_0) = I_0(z_t, \Omega; \Omega_0) \exp \left[-4\gamma q \frac{u_0(\mu)u_0(\mu_0)}{I_0(z_t, \Omega; \Omega_0)} \right]. \quad (23)$$

Here $I_0(z_t, \Omega, \Omega_0)$ is the exiting radiance in the case of $\omega_0 = 1$, $\gamma = \sqrt{3(1-g)(1-\omega_0)}$, $q = \frac{1}{3(1-g)}$ and $u_0(\mu_0) = (1 + \frac{3}{2}\mu_0)/2$. If we substitute into (23) the approximate values of the parameters ($g = 0.86$, $u_0(\mu_0) \approx 1$, $\mu_0 \approx 1$, $I_0(z_t, \Omega, \Omega_0) \approx 0.5$), we find that I depends on ω_0 as $\exp(-10\sqrt{(1-\omega_0)})$. Therefore, the dependence of ΔI on $\Delta\omega_0$ could be linear for $\tau \gg 1$ only when $\Delta\omega_0 \ll 1 - \omega_0$, and that is why ($1 - \omega_0 = 0.002$) the relative error at $|\Delta\omega_0| > 0.01$ in Table 7 is so high. The perturbation technique points to it indirectly by giving the very high value for $dI/d\omega_0$, which is approximately ten times greater than the exiting radiance.

Let us now look closely at Tables 8 and 9. They contain the results for the important case of small cloud contamination, which is difficult to detect with a high level of confidence. The tables show that the perturbation technique allows the effect to be predicted with the above-mentioned accuracy, except in the case of very large angles of illumination and observation. It is not surprising that the error of the prediction is less for model II than for model I at the same optical thickness of the cloud component because of the optical thickness of the lowermost layer of the model II is five times greater than for model I. However, despite such a difference, the sensitivity of the exiting radiance to a cloud contamination is substantially higher for model II than for model I.

5. Conclusion

The above comparison shows that the theoretically established relationship (12) between a small variation of the atmospheric parameters and the corresponding changes in the exiting radiance can be used for both clear and polluted aerosol atmospheres, even with the presence of small cloud contamination. Moreover, it can be the basis of a retrieval algorithm. Let us consider the basic idea of the MISR retrieval algorithm. The aerosol atmosphere is considered as

a stratified medium, whose each layer can be contributed to by 11 possible basic aerosols (each model has its own range of altitudes). Starting from some initial model and following a sophisticated technique, based on using a pre-calculated radiance database, the aerosol optical thickness is retrieved. The perturbation technique can be incorporated into this algorithm in a natural way. Let us introduce a vector $\vec{T} = (\tau_1, \dots, \tau_{11})$ (τ_k is the optical thickness of the k th pure component), which characterizes our aerosol atmosphere model. We also have the measured radiances, I_M , by satellite sensors, and the estimated ones, I_E , for the same geometry, but for some model \vec{T}_0 . The perturbation technique provides the necessary correction, $\Delta\vec{T}$, to the model \vec{T}_0 , on the basis of the comparison between I_M and I_E , which has the form

$$\Delta I = I_M - I_E = \frac{\partial I}{\partial \vec{T}_0} \Delta \vec{T}. \quad (23)$$

Note that the matrix $\frac{\partial I}{\partial \vec{T}_0}$ is calculated together with I_E without noticeable addition to the computer time. It is difficult to estimate for the very first pixel how many iterations it takes to get the final solution, because of it strongly depends on the quality of the initial model. However, when the retrieval is started for the neighboring pixels, the situation is changed. Taking into account that the aerosol profiles at the neighboring pixels differ slightly as a rule, a few iterations should be enough to get the final solution. Additionally, equation (23) also allows us to obtain natural estimation of the retrieval errors for the each pure component if the calibration error is known.

However, the main goal of this paper is to show the applicability of the perturbation technique to describe how the variation of the aerosol atmosphere characteristics effects the exiting radiance for realistic atmosphere models. The comparison of the perturbation technique predictions with the results of the direct solution of the RTE shows its high accuracy. We underline that to use the perturbation technique we have to solve the RTE only once, plus some extra calculation. Note that if we need to analyze the dependence on a single parameter the time

costs of both the perturbation technique and direct simulation are more or less similar, but if we need to investigate multi-parameters dependences the advantages of the perturbation techniques are incontestable. That is why the perturbation technique can be successfully used to make sensitivity analyses of a remote sensing experiment on the basis of consideration of very complex models and, as mentioned above, as a core of the retrieval algorithm. These topics have not been covered in this paper, but will be considered in future work.

6. Acknowledgments

This work is supported by the Australian Research Council.

References

- Bell, G.I., and S. Glasstone, 1970: *Nuclear Reactor Theory*, van Nostrand Reinhold, New York, 619 pp.
- Bates, T.S., B.J. Huebert, J.L. Gras, F.B. Griffiths and P.A. Durkee, 1998: International Global Atmospheric Chemistry (IGAS) Project's First Aerosol Characterisation Experiment (ACE 1), *J. Geophys. Res.*, **103**, 16,297-16,318
- Box, M.A., S.A.W.Gerstl and C.Simmer, 1988a: Application of the adjoint formulation to the calculation of atmospheric radiative effects, *Beitr.Phys. Atmosph.*, **61**, 303-311.
- Box, M.A., M.Keevers and B.H.J. McKellar, 1988b: On the perturbation series for radiative effects, *J.Quant. Spectrosc. Radiat. Transfer*, **39**, 219-223.
- Box, M.A., S.A.W.Gerstl and C.Simmer, 1989a: Computation of atmospheric radiative effects via perturbation theory, *Beitr.Phys. Atmosph.*, **62**, 193-199.
- Box, M.A., B.Croke, S.A.W.Gerstl and C.Simmer, 1989b: Application of the perturbation theory for atmospheric radiative effects: aerosol scattering atmospheres, *Beitr.Phys. Atmosph.*, **62**, 200-211.
- Box, M.A., and C.Sendra, 1995: Sensitivity of exiting radiances to details of the scattering phase function, *J.Quant. Spectrosc. Radiat. Transfer*, **54**, 695-703.
- Charlson, R.J., S.E. Schwartz, J.M. Hales, R.D. Cess, J.A. Coakely, Jr., J.E. Hansen, and D.J. Hofmann, 1992: Climate forcing by antropogenic erosols, *Science*, **255**, 423-430.
- Curry, J.A., 1995: Interactions among aerosols, clouds, and climate of the Arctic Ocean, *The Science of the Total Environment*, **160-161**, 777-791.
- d'Almeida, G.A., P.Koepke, E.P. Shette, 1991: *Atmospheric Aerosols: Global Climatology and Radiative Characteristics*, A.Deepak, Hampton, Va., 559 pp.
- Dave, J.V., 1975: A direct solution of the spherical harmonics approximation to the radiative equation for an arbitrary solar elevation. Part I: Theory, *J.Atmos.Sci.*, **32**, 790 – 798.

- Deschamps, P-Y., F-M. Breon, M. Leroy, A. Podaire, A. Brickaud, J-C. Buriez and G. Seze, 1994: The POLDER mission: Instrument characteristics and scientific objectives, *IEEE*, **32**, 598-615.
- Deirmendjan, D., 1969: *Electromagnetic scattering on spherical polydispersion*, Elsevier, New York, 290 pp.
- Diner, D.J. and C.J. Bruegge, J.V. Martonchik, G.W. Bothwell, E.D. Danielson, V.G. Ford, L.E. Hovland, K.L. Jones, M.L. White (1991). A Multi-angle Imaging SpectroRadiometer for terrestrial remote sensing from the Earth Observing System. *Internatl. J. Imaging Sys. and Tech.*, **3**, 92-107.
- Fletcher, R., 1987: *Practical methods of optimization*, Wiley, Chichester, 436p.
- Gerstl, S.A.W., 1980: Application of modern neutron transport methods to atmospheric radiative transfer, Volume of extended abstracts, International Radiation Symposium, Fort-Collins, Colorado, 11-16 August, 500-502.
- Huebert, B.J., A.Pszenny and B. Blomquist, 1996: The ASTEX/MAGE experiment, *J. Geophys. Res.*, **101**, 4319-4329.
- Marchuk, G.I., 1964: Equation for the value of information from weather satellite and formulation of inverse problems, *Cosmic Res*, **2**, 394-409.
- Kaufman, Y. J., D. Tanre, H.R. Gordon, T.Nakajima, J.Lenoble, R.Frouin, H. Grasll, B.M. Herman, M.D. King, and P.M.Teillet, 1997: Passive remote sensing of tropospheric aerosol and atmosphere correction for the aerosol effect, *J. Geophys. Res.*, **102**, 16,815-16,830.
- Kaufman, Y. J., P.V. Hobbs, V.W.J.H.Kirchhoff, P. Artaxo, L.A. Remer, B.N. Holben, M.D. King, D.E. Ward, E.M. Prins, K.M. Longo, L.F. Mattos, C.A. Nobre, J.D. Spinhirne, Q. Ji, A.M. Thompson, J.F. Gleason, S.A. Christopher and S.-C. Tsay, 1998: Smoke, Clouds, and Radiation-Brazil (SCAR-B) experiment, *J. Geophys. Res.*, **103**, 31,783-31,808
- Lenoble, J., (ed.), 1985: *Radiative Transfer in scattering and absorbing atmospheres: standard computational procedures*, A.Deepak Publishing, Hampton, Va., USA, 300 pp.

- Lewins, J.L., 1965: *Importance: The adjoint function*, Pergamon, New-York, 172 pp.
- Martonchik J.V., D.J. Diner, R.A. Kahn, T.P. Ackerman, M.M. Verstraete, B.Pinty and H.R. Gordon, 1998, *IEEE Trans. On GeoSci. And Remote Sensing*, **36**, 1212-1227.
- Mishchenko, M. I., and L.D. Travis, 1997a: Satellite retrieval of aerosol properties over the ocean using polarization as well as intensity of reflected sunlight, *J. Geophys. Res.*, **102**, 16,989-17,014.
- Mishchenko, M. I., and L.D. Travis, 1997b: Satellite retrieval of aerosol properties over the ocean using measurements of reflected sunlight: Effect of instrumental errors and aerosol absorption, *J. Geophys. Res.*, **102**, 13,543-13,554.
- Rao, C.R.N., L.L Stowe and E.P.McClain, 1989: Remote sensing of aerosols over the oceans using AVHRR data. Theory, practice and applications, *Int. J. Remote Sensing*, **10**, 743-749.
- Russell, P.B., P.V. Hobbs and L.L. Stowe, 1999: Aerosol properties and radiative effects in the United States East Coast haze plume: an overview of the Tropospheric Aerosol Radiative Forcing Observational Experiment (TARFOX), *J. Geophys. Res.*, **104**, 2213-2222.
- Sendra C. and M.A. Box, 2000: Retrieval of the phase function and scattering optical thickness of aerosols: a radiative perturbation theory application, *J.Quant. Spectrosc. Radiat. Transfer*, **64**, 499-515.
- Trautmann, T., P.J. Elliott and M.A. Box, 1992: Shortwave radiative effects of standard aerosol models: a perturbation approach, *Beitr.Phys. Atmosph.*, **65**, 59-78.
- Ustinov, E.A., 1990: The inverse thermal problem: to recover the vertical profile of the absorption coefficient of an optically active component of a planetary atmosphere from observing the emitted radiation, *Cosmic Res*, **28**, 347-355.
- Ustinov, E.A., 1991a: The inverse problem of the photometry of solar radiation reflected by an optically thick planetary atmosphere. Mathematical methods and weighting functions of linearized inverse problem, *Cosmic Res*, **29**, 519-532.

- Ustinov, E.A., 1991b: The inverse problem of the photometry of solar radiation reflected by an optically thick planetary atmosphere. 2. Numerical aspects and requirements on the observation geometry, *Cosmic Res*, **29**, 785-800.
- Ustinov, E.A., 1992: The inverse problem of the photometry of solar radiation reflected by an optically thick planetary atmosphere. 3. Remote sensing of minor gaseous constituents and an atmospheric aerosol, *Cosmic Res*, **30**, 170-181.
- Zege E.P., A.P. Ivanov, and I.L. Katsev, 1991: *Image Transfer through a Scattering Medium*, Springer-Verlag, 360 pp.

Table 1 Atmosphere model.

Layer No.	Altitude, [km]	Molecular atmosphere	Type I	Type II	Type III
1	0-2	(0.0212)	Continental (0.2)	Urban (1.0)	Continental (0.2) Cloud C.1 (10)
2	2-12	(0.0585)	Continental (0.02)		
3	12-30	(0.0178)	Stratospheric (0.003)		
4	30-100	(0.0012)	Upper atmosphere (0)		

The number in brackets are the optical thickness of the component; the aerosol models correspond to the WMO standard, the cloud model is from (Deirmendjan, 1969) .

Table 2. Comparison of the perturbation calculation of $\frac{dI}{d\sigma_e}$ due to the variation of the extinction coefficient of the lowermost layer of Type I atmosphere with results of direct calculation $\frac{\Delta I}{\Delta\sigma_e}$ at variation of extinction coefficient on $\Delta\sigma_e$.

μ_S	μ_R	I	$\frac{dI}{d\sigma_e}$	$\frac{\Delta I}{\Delta\sigma_e}$					
				$\Delta\sigma_e$					
				$0.01\sigma_e$	$0.1\sigma_e$	$1.0\sigma_e$	$-0.05\sigma_e$	$-0.1\sigma_e$	$-0.5\sigma_e$
1.0	1.0	0.9176E-01	0.1852E-01	0.1851E-01	0.1847E-01	0.1808E-01	0.1854E-01	0.1856E-01	0.1876E-01
		relative error		0.000	0.002	0.024	-0.001	-0.003	-0.013
1.0	0.8	0.8997E-01	0.1782E-01	0.1782E-01	0.1786E-01	0.1811E-01	0.1780E-01	0.1777E-01	0.1756E-01
		relative error		0.000	-0.002	-0.016	0.001	0.003	0.015
1.0	0.6	0.9669E-01	0.2215E-01	0.2215E-01	0.2218E-01	0.2226E-01	0.2212E-01	0.2210E-01	0.2184E-01
		relative error		0.000	-0.002	-0.005	0.001	0.002	0.014
1.0	0.2	0.1517E+00	0.3722E-01	0.3714E-01	0.3643E-01	0.3034E-01	0.3763E-01	0.3804E-01	0.4155E-01
		relative error		0.002	0.022	0.227	-0.011	-0.022	-0.104
0.8	0.8	0.7674E-01	0.1767E-01	0.1767E-01	0.1772E-01	0.1797E-01	0.1763E-01	0.1761E-01	0.1732E-01
		relative error		0.000	-0.002	-0.016	0.002	0.004	0.020
0.8	0.6	0.8653E-01	0.2360E-01	0.2360E-01	0.2362E-01	0.2345E-01	0.2358E-01	0.2356E-01	0.2333E-01
		relative error		0.000	-0.001	0.006	0.001	0.002	0.011
0.8	0.2	0.1491E+00	0.4131E-01	0.4122E-01	0.4038E-01	0.3323E-01	0.4179E-01	0.4227E-01	0.4637E-01
		relative error		0.002	0.023	0.243	-0.011	-0.023	-0.109
0.6	0.6	0.7676E-01	0.2450E-01	0.2449E-01	0.2445E-01	0.2364E-01	0.2451E-01	0.2453E-01	0.2454E-01
		relative error		0.000	0.002	0.036	-0.001	-0.001	-0.002
0.6	0.2	0.1451E+00	0.4373E-01	0.4362E-01	0.4263E-01	0.3426E-01	0.4430E-01	0.4487E-01	0.4976E-01
		relative error		0.002	0.026	0.276	-0.013	-0.025	-0.121
0.2	0.2	0.1102E+00	0.2640E-01	0.2628E-01	0.2517E-01	0.1715E-01	0.2705E-01	0.2773E-01	0.3402E-01
		relative error		0.005	0.049	0.539	-0.024	-0.048	-0.224

Table 3. The same as Table 2, but for Type II atmosphere.

μ_S	μ_R	I	$\frac{dI}{d\sigma_e}$	$\frac{\Delta I}{\Delta\sigma_e}$												
				$\Delta\sigma_e$												
				$0.01\sigma_e$	$0.1\sigma_e$	$1.0\sigma_e$	$-0.05\sigma_e$	$-0.1\sigma_e$	$-0.5\sigma_e$							
1.0	1.0	0.9058E-01	0.1066E-01	0.1061E-01	0.1017E-01	0.6812E-02	0.1092E-01	0.1118E-01	0.1361E-01	relative error	0.005	0.048	0.565	-0.023	-0.046	-0.217
1.0	0.8	0.8681E-01	0.1041E-01	0.1037E-01	0.9970E-02	0.6656E-02	0.1063E-01	0.1086E-01	0.1262E-01	relative error	0.004	0.044	0.564	-0.021	-0.041	-0.175
1.0	0.6	0.9439E-01	0.1204E-01	0.1197E-01	0.1135E-01	0.6837E-02	0.1240E-01	0.1276E-01	0.1589E-01	relative error	0.006	0.060	0.761	-0.029	-0.057	-0.242
1.0	0.2	0.1395E+00	0.6678E-02	0.6612E-02	0.6069E-02	0.3197E-02	0.7026E-02	0.7410E-02	0.1260E-01	relative error	0.010	0.100	1.089	-0.050	-0.099	-0.470
0.8	0.8	0.7601E-01	0.1074E-01	0.1068E-01	0.1017E-01	0.6277E-02	0.1103E-01	0.1133E-01	0.1381E-01	relative error	0.005	0.056	0.711	-0.027	-0.052	-0.222
0.8	0.6	0.8781E-01	0.1283E-01	0.1274E-01	0.1194E-01	0.6629E-02	0.1330E-01	0.1379E-01	0.1829E-01	relative error	0.007	0.074	0.935	-0.035	-0.069	-0.298
0.8	0.2	0.1402E+00	0.7308E-02	0.7227E-02	0.6564E-02	0.3197E-02	0.7736E-02	0.8208E-02	0.1472E-01	relative error	0.011	0.113	1.286	-0.055	-0.110	-0.504
0.6	0.6	0.7992E-01	0.1148E-01	0.1138E-01	0.1051E-01	0.5285E-02	0.1201E-01	0.1256E-01	0.1817E-01	relative error	0.009	0.092	1.172	-0.044	-0.086	-0.368
0.6	0.2	0.1378E+00	0.6622E-02	0.6538E-02	0.5854E-02	0.2599E-02	0.7069E-02	0.7566E-02	0.1473E-01	relative error	0.013	0.131	1.548	-0.063	-0.125	-0.550
0.2	0.2	0.9966E-01	0.1245E-02	0.1224E-02	0.1061E-02	0.4171E-03	0.1358E-02	0.1489E-02	0.4056E-02	relative error	0.017	0.173	1.984	-0.083	-0.164	-0.693

Table 4. The same as Table 2, but for Type III atmosphere.

μ_S	μ_R	I	$\frac{dI}{d\sigma_e}$	$\frac{\Delta I}{\Delta\sigma_e}$					
				$\Delta\sigma_e$					
				$0.01\sigma_e$	$0.1\sigma_e$	$1.0\sigma_e$	$-0.05\sigma_e$	$-0.1\sigma_e$	$-0.5\sigma_e$
1.0	1.0	0.5165E+00	0.2918E+00	.2905E+00 0.005	0.2787E+00 0.047	0.1896E+00 0.539	0.2986E+00 -0.023	0.3055E+00 -0.045	0.3596E+00 -0.188
1.0	0.8	0.4887E+00	0.2619E+00	.2605E+00 0.005	0.2488E+00 0.053	0.1665E+00 0.573	0.2688E+00 -0.026	0.2759E+00 -0.051	0.3404E+00 -0.231
1.0	0.6	0.4402E+00	0.2241E+00	.2229E+00 0.005	0.2125E+00 0.054	0.1415E+00 0.584	0.2303E+00 -0.027	0.2368E+00 -0.053	0.3000E+00 -0.253
1.0	0.2	0.3721E+00	0.1392E+00	.1385E+00 0.005	0.1321E+00 0.054	0.8800E-01 0.582	0.1430E+00 -0.027	0.1470E+00 -0.053	0.1859E+00 -0.251
0.8	0.8	0.3826E+00	0.1881E+00	.1870E+00 0.006	0.1778E+00 0.058	0.1169E+00 0.609	0.1937E+00 -0.029	0.1996E+00 -0.057	0.2598E+00 -0.276
0.8	0.6	0.3920E+00	0.1610E+00	.1601E+00 0.006	0.1519E+00 0.060	0.9939E-01 0.620	0.1660E+00 -0.030	0.1713E+00 -0.060	0.2299E+00 -0.300
0.8	0.2	0.3614E+00	0.1000E+00	.9943E-01 0.006	0.9438E-01 0.060	0.6181E-01 0.618	0.1031E+00 -0.030	0.1064E+00 -0.060	0.1425E+00 -0.298
0.6	0.6	0.3186E+00	0.1034E+00	.1028E+00 0.006	0.9734E-01 0.062	0.6336E-01 0.632	0.1067E+00 -0.031	0.1103E+00 -0.063	0.1530E+00 -0.324
0.6	0.2	0.3295E+00	0.6422E-01	.6383E-01 0.006	0.6048E-01 0.062	0.3940E-01 0.630	0.6628E-01 -0.031	0.6848E-01 -0.062	0.9482E-01 -0.323
0.2	0.2	0.1673E+00	0.1330E-01	.1322E-01 0.006	0.1253E-01 0.061	0.8169E-02 0.628	0.1372E-01 -0.031	0.1417E-01 -0.062	0.1958E-01 -0.321

Table 5. Comparison of the perturbation calculation of $\frac{dI}{d\omega_0}$ due to the variation of the single scattering albedo of the

lowermost layer of Type I model atmosphere with results of direct calculation $\frac{\Delta I}{\Delta\omega_0}$ at the single scattering albedo on $\Delta\omega_0$.

μ_S	μ_R	I	$\frac{dI}{d\omega_0}$	$\frac{\Delta I}{\Delta\omega_0}$					
				$\Delta\omega_0$					
				0.01	0.05	-0.01	-0.05	-0.1	-0.5
1.0	1.0	0.9176E-01	0.5080E-01	0.5097E-01	0.5166E-01	0.5063E-01	0.4997E-01	0.4917E-01	0.4376E-01
		relative error		-0.003	-0.017	0.003	0.017	0.033	0.161
1.0	0.8	0.8997E-01	0.5404E-01	0.5427E-01	0.5521E-01	0.5381E-01	0.5292E-01	0.5184E-01	0.4463E-01
		relative error		-0.004	-0.021	0.004	0.021	0.042	0.211
1.0	0.6	0.9669E-01	0.6676E-01	0.6709E-01	0.6843E-01	0.6644E-01	0.6517E-01	0.6365E-01	0.5352E-01
		relative error		-0.005	-0.024	0.005	0.024	0.049	0.247
1.0	0.2	0.1517E+00	0.1259E+00	0.1267E+00	0.1296E+00	0.1252E+00	0.1225E+00	0.1192E+00	0.9732E-01
		relative error		-0.006	-0.028	0.006	0.028	0.057	0.294
0.8	0.8	0.7674E-01	0.5184E-01	0.5208E-01	0.5311E-01	0.5158E-01	0.5060E-01	0.4943E-01	0.4165E-01
		relative error		-0.005	-0.024	0.005	0.024	0.049	0.245
0.8	0.6	0.8653E-01	0.6655E-01	0.6690E-01	0.6837E-01	0.6618E-01	0.6480E-01	0.6313E-01	0.5215E-01
		relative error		-0.005	-0.027	0.005	0.027	0.054	0.276
0.8	0.2	0.1491E+00	0.1301E+00	0.1309E+00	0.1341E+00	0.1293E+00	0.1263E+00	0.1227E+00	0.9897E-01
		relative error		-0.006	-0.030	0.006	0.030	0.060	0.314
0.6	0.6	0.7676E-01	0.6573E-01	0.6610E-01	0.6768E-01	0.6534E-01	0.6386E-01	0.6208E-01	0.5045E-01
		relative error		-0.006	-0.029	0.006	0.029	0.059	0.303
0.6	0.2	0.1451E+00	0.1320E+00	0.1329E+00	0.1363E+00	0.1312E+00	0.1280E+00	0.1241E+00	0.9897E-01
		relative error		-0.006	-0.031	0.006	0.032	0.064	0.334
0.2	0.2	0.1102E+00	0.9256E-01	0.9318E-01	0.9570E-01	0.9196E-01	0.8959E-01	0.8677E-01	0.6853E-01
		relative error		-0.007	-0.033	0.007	0.033	0.067	0.351

Table 6. The same as Table 2, but for Type II atmosphere.

μ_S	μ_R	I	$\frac{dI}{d\omega_0}$	$\frac{\Delta I}{\Delta\omega_0}$						
				$\Delta\omega_0$						
				0.01	0.05	0.1	-0.01	-0.05	-0.1	-0.5
1.0	1.0	0.9058E-01	0.1565E+00	0.1583E+00	0.1660E+00	0.1767E+00	0.1547E+00	0.1480E+00	0.1403E+00	0.9954E-01
		relative error		-0.011	-0.057	-0.115	0.011	0.057	0.115	0.572
1.0	0.8	0.8681E-01	0.1659E+00	0.1681E+00	0.1776E+00	0.1909E+00	0.1637E+00	0.1554E+00	0.1461E+00	0.9690E-01
		relative error		-0.013	-0.066	-0.131	0.013	0.067	0.135	0.712
1.0	0.6	0.9439E-01	0.1914E+00	0.1942E+00	0.2058E+00	0.2221E+00	0.1888E+00	0.1787E+00	0.1673E+00	0.1079E+00
		relative error		-0.014	-0.070	-0.138	0.014	0.071	0.144	0.773
1.0	0.2	0.1395E+00	0.2194E+00	0.2224E+00	0.2352E+00	0.2531E+00	0.2165E+00	0.2054E+00	0.1929E+00	0.1277E+00
		relative error		-0.014	-0.067	-0.133	0.014	0.068	0.137	0.718
0.8	0.8	0.7601E-01	0.1556E+00	0.1579E+00	0.1673E+00	0.1806E+00	0.1534E+00	0.1452E+00	0.1360E+00	0.8761E-01
		relative error		-0.014	-0.070	-0.138	0.014	0.072	0.145	0.776
0.8	0.6	0.8781E-01	0.1861E+00	0.1889E+00	0.2006E+00	0.2170E+00	0.1834E+00	0.1733E+00	0.1619E+00	0.1027E+00
		relative error		-0.015	-0.072	-0.142	0.015	0.074	0.150	0.812
0.8	0.2	0.1402E+00	0.2238E+00	0.2269E+00	0.2401E+00	0.2586E+00	0.2208E+00	0.2093E+00	0.1965E+00	0.1293E+00
		relative error		-0.014	-0.068	-0.135	0.014	0.069	0.139	0.730
0.6	0.6	0.7992E-01	0.1714E+00	0.1739E+00	0.1849E+00	0.2002E+00	0.1688E+00	0.1594E+00	0.1488E+00	0.9374E-01
		relative error		-0.015	-0.073	-0.144	0.015	0.075	0.152	0.828
0.6	0.2	0.1378E+00	0.2172E+00	0.2202E+00	0.2329E+00	0.2507E+00	0.2143E+00	0.2033E+00	0.1909E+00	0.1259E+00
		relative error		-0.014	-0.067	-0.134	0.014	0.068	0.138	0.726
0.2	0.2	0.9966E-01	0.1120E+00	0.1134E+00	0.1190E+00	0.1267E+00	0.1108E+00	0.1059E+00	0.1003E+00	0.7002E-01
		relative error		-0.012	-0.058	-0.116	0.012	0.058	0.117	0.600

Table 7. The same as Table 2, but for Type III atmosphere.

μ_S	μ_R	I	$\frac{dI}{d\omega_0}$	$\frac{\Delta I}{\Delta\omega_0}$			
				$\Delta\omega_0$			
				-0.01	-0.05	-0.1	-0.5
1.0	1.0	0.5165E+00	0.5129E+01	0.4665E+01	0.3416E+01	0.2552E+01	0.8573E+00
		relative error		0.099	0.501	1.010	4.982
1.0	0.8	0.4887E+00	0.5310E+01	0.4824E+01	0.3514E+01	0.2609E+01	0.8402E+00
		relative error		0.101	0.511	1.035	5.320
1.0	0.6	0.4402E+00	0.5170E+01	0.4691E+01	0.3404E+01	0.2516E+01	0.7800E+00
		relative error		0.102	0.519	1.055	5.628
1.0	0.2	0.3721E+00	0.3701E+01	0.3359E+01	0.2441E+01	0.1809E+01	0.5738E+00
		relative error		0.102	0.516	1.046	5.450
0.8	0.8	0.3826E+00	0.4483E+01	0.4070E+01	0.2961E+01	0.2193E+01	0.6826E+00
		relative error		0.101	0.514	1.044	5.567
0.8	0.6	0.3920E+00	0.4484E+01	0.4076E+01	0.2976E+01	0.2212E+01	0.6938E+00
		relative error		0.100	0.507	1.027	5.463
0.8	0.2	0.3614E+00	0.3317E+01	0.3023E+01	0.2228E+01	0.1673E+01	0.5507E+00
		relative error		0.097	0.489	0.982	5.022
0.6	0.6	0.3186E+00	0.3443E+01	0.3137E+01	0.2310E+01	0.1732E+01	0.5560E+00
		relative error		0.097	0.490	0.988	5.192
0.6	0.2	0.3295E+00	0.2659E+01	0.2437E+01	0.1828E+01	0.1397E+01	0.4855E+00
		relative error		0.091	0.454	0.903	4.478
0.2	0.2	0.1673E+00	0.7810E+00	0.7246E+00	0.5687E+00	0.4545E+00	0.1867E+00
		relative error		0.078	0.373	0.719	3.184

Table 8. Comparison of perturbation calculation of $\frac{dI}{d\tau_c}$ with results of direct calculation $\frac{\Delta I}{\Delta\tau_c}$ at adding the cloud optical thickness $\Delta\tau_c$ to the lowermost layer of Type I model atmosphere.

μ_S	μ_R	I	$\frac{dI}{d\tau_c}$	$\frac{\Delta I}{\Delta\tau_c}$			
				$\Delta\tau_c$			
				0.01	0.05	0.1	0.5
1.0	1.0	0.4990E-01	0.3695E-01	0.3675E-01	0.3598E-01	0.3507E-01	0.2959E-01
		relative error		0.0054	0.0270	0.0536	0.2485
1.0	0.8	0.4930E-01	0.2724E-01	0.2714E-01	0.2675E-01	0.2630E-01	0.2375E-01
		relative error		0.0037	0.0183	0.0357	0.1470
1.0	0.6	0.5802E-01	0.1403E-01	0.1409E-01	0.1430E-01	0.1456E-01	0.1621E-01
		relative error		-0.0039	-0.0190	-0.0366	-0.1346
1.0	0.2	0.1243E+00	0.1807E-01	0.1791E-01	0.1735E-01	0.1677E-01	0.1435E-01
		relative error		0.0088	0.0413	0.0774	0.2593
0.8	0.8	0.4226E-01	0.1029E-01	0.1032E-01	0.1065E-01	0.1104E-01	0.1355E-01
		relative error		-0.0027	-0.0336	-0.0683	-0.2409
0.8	0.6	0.6175E-01	0.1926E-01	0.1925E-01	0.1974E-01	0.2029E-01	0.2287E-01
		relative error		0.0009	-0.0239	-0.0506	-0.1577
0.8	0.2	0.2012E+00	0.6045E-01	0.5954E-01	0.5746E-01	0.5492E-01	0.3951E-01
		relative error		0.0153	0.0521	0.1007	0.5301
0.6	0.6	0.7811E-01	0.3244E-01	0.3231E-01	0.3290E-01	0.3346E-01	0.3398E-01
		relative error		0.0040	-0.0138	-0.0306	-0.0454
0.6	0.2	0.2986E+00	0.1360E+00	0.1335E+00	0.1272E+00	0.1196E+00	0.7690E-01
		relative error		0.0191	0.0694	0.1368	0.7686
0.2	0.2	0.4743E+00	0.2750E+00	0.2663E+00	0.2401E+00	0.2125E+00	0.1042E+00
		relative error		0.0324	0.1452	0.2941	1.6396

Table 9. The same as Table 8, but for Type II model atmosphere.

μ_S	μ_R	I	$\frac{dI}{d\tau_C}$	$\frac{\Delta I}{\Delta\tau_C}$			
				$\Delta\tau_C$			
				0.01	0.05	0.1	0.5
1.0	1.0	0.7410E-01	0.1033E+00	0.1029E+00	0.1015E+00	0.9986E-01	0.8913E-01
		relative error		0.0035	0.0172	0.0341	0.1585
1.0	0.8	0.7266E-01	0.7840E-01	0.7821E-01	0.7748E-01	0.7660E-01	0.7093E-01
		relative error		0.0024	0.0119	0.0235	0.1054
1.0	0.6	0.8317E-01	0.4320E-01	0.4324E-01	0.4341E-01	0.4360E-01	0.4460E-01
		relative error		-0.0010	-0.0049	-0.0093	-0.0314
1.0	0.2	0.1346E+00	0.2782E-01	0.2782E-01	0.2785E-01	0.2788E-01	0.2808E-01
		relative error		-0.0002	-0.0011	-0.0023	-0.0093
0.8	0.8	0.6709E-01	0.3442E-01	0.3504E-01	0.3530E-01	0.3562E-01	0.3752E-01
		relative error		-0.0176	-0.0250	-0.0336	-0.0826
0.8	0.6	0.9445E-01	0.4791E-01	0.4948E-01	0.4965E-01	0.4983E-01	0.5042E-01
		relative error		-0.0317	-0.0350	-0.0386	-0.0497
0.8	0.2	0.2157E+00	0.6448E-01	0.6954E-01	0.6892E-01	0.6816E-01	0.6265E-01
		relative error		-0.0727	-0.0644	-0.0540	0.0293
0.6	0.6	0.1131E+00	0.6182E-01	0.6441E-01	0.6435E-01	0.6423E-01	0.6218E-01
		relative error		-0.0403	-0.0394	-0.0376	-0.0058
0.6	0.2	0.3081E+00	0.1430E+00	0.1498E+00	0.1474E+00	0.1445E+00	0.1249E+00
		relative error		-0.0454	-0.0298	-0.0104	0.1448
0.2	0.2	0.4281E+00	0.2851E+00	0.2879E+00	0.2802E+00	0.2711E+00	0.2154E+00
		relative error		-0.0096	0.0178	0.0519	0.3235