

**Development of a Global Validation Package for Satellite Oceanic Aerosol Optical  
Thickness Retrieval Based on AERONET Observations and Its Application to  
NOAA/NESDIS Operational Aerosol Retrievals**

Tom X.-P. Zhao<sup>1</sup>, Larry L. Stowe<sup>2</sup>, Alexander Smirnov<sup>3</sup>, David Crosby<sup>2</sup>, John Sapper<sup>4</sup>,  
and Charles R. McClain<sup>5</sup>

<sup>1</sup>*CIRA Visiting Scientist, NOAA/NESDIS/ORA, Camp Springs, MD 20746*

<sup>2</sup>*NOAA/NESDIS/ORA, Camp Spring, MD 20746*

<sup>3</sup>*NASA/GSFC, SSAI, Code 923, Greenbelt, MD 20771*

<sup>4</sup>*NOAA/NESDIS, OSDPD, Suitland, Washington D. C. 20395*

<sup>5</sup>*NASA/GSFC, Code 970.2, Greenbelt, MD 20771*

Corresponding author and address: Dr. Tom X-P. Zhao  
E/RA1, RM 711-C, WWBG, NOAA/NESDIS/ORA  
5200 Auth Road, Camp Springs, MD 20746-4304  
Phone: (301)763-8059 (ext. 213)  
Fax: (301)763-8108  
Email: [xzhao@nesdis.noaa.gov](mailto:xzhao@nesdis.noaa.gov)

**Submitted to the GLOBAL AEROSOL CLIMATOLOGY special issue of *Journal***

## ABSTRACT

In this paper, a global validation package for satellite aerosol optical thickness retrieval using the Aerosol Robotic Network (AERONET) observations as ground truth is described. To standardize the validation procedure, the optimum time/space match-up window, the ensemble statistical analysis method, the best selection of AERONET channels and the numerical scheme used to interpolate/extrapolate these observations to satellite channels have been identified through sensitivity studies. The package is shown to be a unique tool for more objective validation and inter-comparison of satellite aerosol retrievals, helping to satisfy an increasingly important requirement of the satellite aerosol remote sensing community. Results of applying the package to the 2<sup>nd</sup> generation operational aerosol observational data (AEROBS) from NOAA-14/AVHRR in 1998 and to the same year aerosol observation data (CERES-SSF4) from TRMM/VIRS are presented as examples of global validation. The usefulness of the package for identifying improvements to the aerosol optical thickness ( $\tau$ ) retrieval algorithm is also demonstrated.

The principal causes of systematic errors in the current NOAA/NESDIS operational aerosol optical thickness retrieval algorithm have been identified and can be reduced significantly if the correction and adjustment suggested from the global validation are

transfer model parameters that reduce systematic errors in  $\tau$  retrievals are suggested for consideration in our next generation algorithm. Basic features that should be included in the next generation algorithm to reduce random error in  $\tau$  retrievals and the resulting error in the effective Angstrom wavelength exponent have also been discussed.

Compared to the AERONET observation, the NOAA-14/AVHRR (AEROBS)  $\tau$  values for mean conditions are biased high by 0.05 and 0.08, with random errors of 0.08 and 0.05, at 0.63 $\mu\text{m}$  and 0.83 $\mu\text{m}$ , respectively. Correspondingly, the TRMM/VIRS (CERES-SSF4) values for mean conditions are biased high by 0.06 and 0.02, with random errors of 0.06 and 0.04, at 0.63 $\mu\text{m}$  and 1.61 $\mu\text{m}$ , respectively. After corrections and adjustments to the retrieval algorithm, the biases in both channels of AVHRR and VIRS are reduced significantly to values close to zero, although random error is almost unchanged. The effective Angstrom wavelength exponent ( $\alpha$ ) derived directly from the aerosol optical thicknesses ( $\tau_s$ ) has been shown to be poorly correlated both before and after adjustments, indicating that random error in the  $\tau$  measurement (possibly related to aerosol model parameter variations or cloud/surface reflectance contamination) needs to be reduced.

## **1. Introduction**

It is generally recognized that the direct and indirect radiative effects of tropospheric aerosols on global climate are comparable to greenhouse gases but with opposite sign and with larger uncertainties (Hansen and Lacis, 1990; Charlson et al., 1992; Lacis and Mishchenko, 1995; IPCC 1996). Long term global aerosol measurement based on satellite aerosol remote sensing may help us to practically reduce these uncertainties (King et al., 1999; Hansen et al., 2000) provided the performance of the retrieval algorithms and instruments is well evaluated and documented. Actually, many validations performed for spaceborne and airborne aerosol retrievals can be found in the literature (e.g., Kaufman et al., 1990; Ignatov et al., 1995a, b; Stowe et al., 1997; Nakajima and Higurashi, 1997; Tanre et al., 1997; Chu et al., 1999; Goloub et al., 1999; Durkee et al., 1999). One common character of these validations is that they were performed only in a limited region and period covered either by an airborne or a ship cruise campaign, or both. Comparable validation results have been obtained from these validations, but with different validation concepts and procedures, which may generate many ambiguities for inter-comparison of these retrieval algorithms.

Lacking global ground-based aerosol observations is the major obstacle for global validation of satellite aerosol retrievals. However, the Aerosol Robotic Network (AERONET [Holben et al., 1998]) initiated by NASA's EOS program, recently

aerosol spectral optical depths, aerosol size distributions, etc., in a manner suitable for integration with satellite data (see e.g., Dubovik et al., 2000, Smirnov et al., 2000).

It is time to standardize procedures for a more objective global validation of satellite aerosol retrievals and their inter-comparison. This is a very important step considering the fact that no one sensor system is capable of providing totally unambiguous information, and hence a careful intercomparison of derived products from different sensors, together with a comprehensive network of ground-based sunphotometer and sky radiometer systems, are required to advance our quantitative understanding of global aerosol characteristics (King et al., 1999).

The 2<sup>nd</sup> generation NOAA/NESDIS operational aerosol optical thickness ( $\tau$ ) retrieval algorithm is used to process data from NOAA-14/AVHRR (Stowe et al., 1997) and TRMM/VIRS (Ignatov and Stowe, 2000) for global oceanic aerosol monitoring. The algorithm provides estimates of  $\tau$  independently in each reflectance channel of AVHRR (0.63 $\mu$ m and 0.83 $\mu$ m) and VIRS (0.63 $\mu$ m and 1.61 $\mu$ m), assuming the molecular atmosphere, the aerosol microphysics and surface reflectance are known and invariant. The effective Angstrom wavelength exponent ( $\alpha$ ) can be derived from the two independent measurements of  $\tau$  in two reflectance channels of AVHRR or VIRS. In operation, the relationship between aerosol optical thickness ( $\tau$ ) and dimensionless reflectance ( $\rho$ , normalized to solar flux) is described by a four-dimensional look-up table

volume scattering and absorption coefficients, and the aerosol phase function derived from Mie calculations with a prescribed aerosol microphysical model. The oceanic albedo is set to 0.002 and 0.0005, and zero for 0.63 $\mu\text{m}$ , 0.83 $\mu\text{m}$ , and 1.61 $\mu\text{m}$  channels, respectively. The aerosol model uses a mono-modal lognormal distribution:  $dN/d \ln r = N(2\mathbf{P})^{-1/2} \ln^{-1}(\mathbf{s}) \exp[-\ln(r/r_m)^2 / 2(\ln \mathbf{s})^2]$ , where mode radius  $r_m=0.1\mu\text{m}$  and variance of size distribution  $\sigma=2.03$ ; a complex index of refraction  $n=1.40-0.0i$  (or albedo of single scattering  $\omega=1$ ). More detailed descriptions on the retrieval algorithm can be found in Stowe et al. (1997).

This algorithm hasn't gone through a complete global validation. The documented performance of the algorithm was only based on three oceanic cruise validations in the tropical and north Atlantic Ocean for its operational one-channel (0.63 $\mu\text{m}$ ) version on NOAA-9 & 11/AVHRR (Ignatov et al., 1995a, b; Stowe et al., 1997). The global aerosol products of AVHRR are to be widely used for studies of radiative forcing on climate change due to its long-term availability, retrospectively back to 1981 in the AVHRR Pathfinder Atmosphere (PATMOS) data (Stowe and Jacobowitz, 1997). Thus, a more complete and objective global validation and error estimation of the retrieval algorithm applied to both channels is necessary.

Based on the above considerations, a global validation package based on AERONET observations has been developed. A prototype for using AERONET data objectively for

aerosol retrieval algorithm) presented, but also the advantages of our global validation package (based on a comprehensive network of ground-based observations) to the improvement of satellite aerosol retrieval algorithms (as anticipated by, for example, King et al., 1999) is demonstrated. Development of the global validation package is described in Section 2. Validation results on the 2<sup>nd</sup> generation NOAA/NESDIS operational aerosol retrieval algorithm, applied to Channel 1 and 2, are given in Section 3. The value of the package for the improvement of an aerosol retrieval algorithm is demonstrated in Section 4. Some important issues are discussed in Section 5. Summary and concluding remarks are given in Section 6.

## **2. Development of the Global Validation Package**

### **a) Basic Concept**

Since aerosol optical thickness  $\tau$  is derived from the measurement of spectral attenuation of the direct solar beam using the CIMEL sun/sky radiometers at AERONET stations (Holben et al., 1998). This measurement is equivalent to sun photometer measurement. Its accuracy is much higher than that derived from backward scattering radiance (contaminated by varying surface (land, ocean, cloud) properties) measured from satellite (Tanre et al., 1996). For convenience, we subsequently refer to these solar extinction measurements as Sun-photometer (SP) data. Thus, AERONET measurement

observations (e.g., Smirnov et al., 2000) are used as ground truth in the validation. Initially, the global validation package was applied to AVHRR and VIRS data from 1998 (the year VIRS data became available) to validate the NESDIS 2<sup>nd</sup> generation retrieval algorithm performance. After checking the Level 2 AERONET SP data, it is found that there aren't Level 2 data for Dakar and Guadeloupe in 1998. There are only two months of Level 2 data in 1998 for Cape Verde (January and February), Ascension Island (November and December), and San Nicolas (November and December). Also, Level 2 data in 1998 at Barbados are missing for some channels. Thus, only 7 of 13 AERONET stations (Andros Island, Bahrain, Bermuda, Dry Tortugas, Kaashidhoo, Surinam, and Lanai, which are highlighted in Table 1) have been kept in our validation for the 1998 satellite observations.

The aerosol optical thickness  $\tau$  and its effective Angstrom wavelength exponent  $\alpha$  retrieved from these two satellite sensors ( $\tau_{st}$  and  $\alpha_{st}$ ) are matched up with the corresponding  $\hat{o}truth\hat{o}$  values derived from the same day surface AERONET SP observations ( $\tau_{sp}$  and  $\alpha_{sp}$ ). They are statistically processed within an optimum space/time window from which scatter diagrams of  $\tau_{st}$  vs.  $\tau_{sp}$  and  $\alpha_{st}$  vs.  $\alpha_{sp}$  are produced. Linear regression analyses are performed, predicting the satellite retrieved values of  $\tau_{st}$  or  $\alpha_{st}$  as a function of the SP values of  $\tau_{sp}$  or  $\alpha_{sp}$  in the form of  $\tau_{st} = A + B\tau_{sp}$  (or  $\alpha_{st} = A + B\alpha_{sp}$ ). Retrieval algorithm performance can be evaluated from resulting statistical parameters of



(proportional error) indicates that there may be some inconsistency between the aerosol microphysical model (such as refractive index) used in the retrieval algorithm and that in the real world. A very good diagnostic analysis of the physical rationale behind the errors represented by non-zero intercept and non-unity slope have been performed in Stowe et al. (1997) using the linearized single scattering approximation of the radiative transfer equation, which is also utilized in the discussion of Section 5.

This validation has been performed not only for single AERONET stations (called regional validation) but also for the ensemble of all selected stations (called global validation hereafter). Since the number of match-up days found for a single AERONET station in 1998 is not sufficient (need probably at least 60 samples) for conclusive regional validation, the paper has been restricted to the global validation.

Our validation package is summarized with a flow chart in Fig.1. Each step (1-12) is briefly described. The original satellite data (AEROBS or CERES-SSF4) and surface AERONET observations (1, 2) are collected and reformatted (3) around the 13 baseline AERONET stations to produce a smaller archived data set (4) for use in subsequent steps. The match-ups are searched (5) according to an optimal time/space window to produce a merged, match-up (archived) data set (6). This archived match-up data set is used to do regression validation (7) and to generate statistical summaries (8). Based on interpretation of these results, conclusions can be published (11, 12). If potential improvements to the

Some important optimizing studies, which are critical to the reliability and consistency of the validation results, are discussed sequentially in the following subsection b).

#### b) Sensitivity Studies and Optimization

The noise in  $\tau_{st}$  and  $\tau_{sp}$  may result not only from their natural variability and measurement errors but also from errors associated with improper statistical treatment of the validation process, in particular, selection of match-up window size and sampling approach. If the validation of a retrieval algorithm is not performed according to an optimal procedure (based on solid quantitative analysis), the resulting performance may not be truly representative. An optimal validation procedure is also very important for the inter-comparison of different satellite aerosol retrieval algorithms since subtle differences in their validation procedures may cause the performance of the algorithms to appear to be different. In other words, if validation procedures are not standardized, one can always adjust the procedures to obtain comparable performance with another algorithm. This is why one sees similar validation performance for different satellite aerosol retrieval algorithms in the literature. It is time to standardize validation procedures for a more objective validation of satellite aerosol retrievals and their inter-comparison. This paper is an attempt to move toward this objective. Initially the importance of aerosol validation

and ground truth) being compared. In our validation operation, daily quality controlled (Level 2) SP data (multi-spectral optical thickness) at AERONET stations are used to compute  $\tau$  and  $\alpha$  for the solar reflectance spectral channels of the satellite instruments (0.63 $\mu\text{m}$ , 0.83 $\mu\text{m}$ , and 1.61 $\mu\text{m}$ ). Error may be introduced by using different schemes to interpolate (or extrapolate, as is the case at 1.61 $\mu\text{m}$ ) aerosol data from SP channels to satellite channels. Sensitivity studies have been performed mainly with respect to these uncertainties, which are discussed below in sequence. These sensitivity studies not only minimize the errors in the validation process but also optimize and standardize the global validation process.

#### 1) Time/Space Match-up Window

Selection of the match-up window is a critical part of the aerosol validation process, since it actually defines the collocation between satellite and sun-photometer observations used in the comparison. We have not found any quantitative analysis of this issue in the aerosol retrieval literature. The representativeness of a validation result without considering this issue carefully would be questionable.

Three AERONET stations with the most Level 2 SP observations in 1998 (Dry Tortugas, Bermuda, and Kaashidhoo) have been chosen as the base stations for this investigation. AVHRR aerosol optical thickness retrievals have been evaluated against SP observations through linear regression analysis ( $\tau_{\text{AVHRR}} = A + B\tau_{\text{SP}}$ ) (or scatter diagrams of

The match-up window of Table 2 consists of an outer circle with variable radial distance from the site, excluding an inner circle (with a fixed radius of 25km) to reduce the effects of coastline or shallow water influences. It is also defined by the time difference (in hours) between the satellite and SP observations. Linear regression coefficients (A, B,  $\sigma$ , and  $R^2$ ) have been derived for all 20 match-up windows listed in Table 2 at the three selected stations. Plots of the resulting regression coefficients at Dry Tortugas are given as an example in Fig.2. Results from the other two sites are similar. The window with intercept (A) closest to zero, slope (B) closest to unity, highest correlation ( $R^2$ ) or lowest standard error ( $\sigma$ ) is considered optimal. If there are any inconsistencies between the coefficients, the highest value of  $R^2$  determines the optimum window. This is because only when the correlation of two different observations is sufficiently large does the relationship between them have meaning. Based on these criteria, it is obvious from Fig.2 that a 1 hour/100km match-up window is optimal for Dry Tortugas.

A similar study with 1 hour time and variable spatial match-up window has been performed for aerosol retrievals from CERES-SSF version 4 data at Bermuda and Kaashidhoo (have most of SSF-SP match-ups in 1998). At Bermuda, the optimal space window is 200km, if judgement is based on the values of coefficients A and B, but it is 100km when judgement is based on the correlation coefficient ( $R^2$ ). Thus, as previously

of AEROBS and SSF-4 data from the  $0.63\mu\text{m}$  channel, respectively. Again, a second order interpolation scheme for the four SP channels ( $0.87\mu\text{m}$ ,  $0.67\mu\text{m}$ ,  $0.50\mu\text{m}$ , and  $0.44\mu\text{m}$ ) has been employed. It is obvious that the regression coefficients are closer to ideal with the 100km match-up window. This shows how critical the selection of an optimal match-up window can be for aerosol validation.

## 2) Interpolation/Extrapolation of SP Observation

The sensitivity study on interpolation/extrapolation of SP wavelength dependent observations to satellite channels ( $0.63\mu\text{m}$ ,  $0.83\mu\text{m}$ , and  $1.61\mu\text{m}$ ) has been performed on two aspects of the problem using the optimal match-up window (1 hour/100km) regression statistics. The first study is on channel selection and the second is on the interpolation scheme. Although sensitivity studies have been performed for both regional and global validations, only results from the global validation, with its larger statistical sample, are presented. Since AERONET SP observations are not available from the same set of channels for all stations, we have examined the validation procedure sensitivity to three selected sets of wavelength channels (listed in Table 4). Channel Set I is considered as the default set, since it is available at almost all selected AERONET stations. Two interpolation schemes, using first order and second order polynomial fits (in natural logarithm of wavelength) to each set of observations, have been selected for testing.

channel data, the 1.02 $\mu\text{m}$  channel is added to SP Channel Set II to form Channel Set III. Sensitivity results with SSF4 data are summarized in Tables 6a and 6b.

In Tables 5a and 5b, the regression coefficients for each AVHRR channel change only slightly by choosing different SP channel sets and interpolation schemes. We have concluded that these small differences are not statistically significant with the following statistical testing approach, using SP Channel Set I along with 1<sup>st</sup> and 2<sup>nd</sup> order interpolation for the 0.63 $\mu\text{m}$  channel as an example.

First, half of the match-up points (e.g., 60 out of 120 for AEROBS) are randomly picked to do one regression analysis and the other half are used to do a second regression analysis. Then, three difference statistics ( $DSP_A$  for intercept A,  $DSP_B$  for slope B, and  $DSP_\sigma$  for standard error) are computed from the following three equations:

$$DSP_A = \frac{A_1 - A_2}{\sqrt{s_{A_1}^2 + s_{A_2}^2}} \quad (1a)$$

$$DSP_B = \frac{B_1 - B_2}{\sqrt{s_{B_1}^2 + s_{B_2}^2}} \quad (1b)$$

$$DSP_s = \frac{s_1^2}{s_2^2} \quad (1c)$$

where,  $A_1$ ,  $B_1$ , and  $\sigma_1$  are intercept, slope, and standard error of the regression line for the first half of match-up points, which uses 1<sup>st</sup> order interpolation of Set I SP data.  $A_2$ ,  $B_2$ , and  $\sigma_2$  are from the regression line for the second half of the match-up points, which uses

standard deviation one. If the errors of the two regressions have the same standard deviation, then  $DSP_{\sigma}$  will have approximately an  $F$  distribution (Ostle and Mensing, 1988; Beyer, 1991). The sample size (60) is large enough to insure that the approximations are reasonable. If  $-1.96 < DSP_A$  (or  $DSP_B$ )  $< +1.96$ , it suggests there is no significant difference (at the 95% confidence level) between the intercept (or slope) using 1<sup>st</sup> and 2<sup>nd</sup> order polynomial interpolation regressions. Similarly, if  $0.60 < DSP_{\sigma} < 1.67$ , it means there is no significant difference (at the 95% confidence level) in the standard error of regression between the 1<sup>st</sup> and 2<sup>nd</sup> order polynomial interpolation schemes. For this example,  $DSP_A = -0.218$ ,  $DSP_B = 0.655$ , and  $DSP_{\sigma} = 0.923$ , supporting the conclusion.

The regression statistics not being sensitive to the selection of SP channel sets or interpolation schemes may be due to the fact that the spectral range covered by all SP channel sets always contains the AVHRR channels (0.63 $\mu$ m and 0.83 $\mu$ m). Thus, only interpolation (no extrapolation) is involved. Since 2<sup>nd</sup> order interpolation has traditionally been used for the validation of satellite aerosol retrievals in our past research (see Ignatov et al., 1995; Stowe et al., 1997) as well as there being no statistically significant differences between the 1<sup>st</sup> and 2<sup>nd</sup> order interpolations, it was decided to continue the use of 2<sup>nd</sup> order interpolation with Set I SP channels for AEROBS validation.

Results of the above sensitivity study applied to TRMM/VIRS (CERES-SSF4) data are presented in Tables 6a and 6b. At 0.63 $\mu$ m, the change of regression statistics for

interpolation scheme. This is probably because Channel 2 of VIRS is beyond the spectral range of the SP channels. Extrapolation is required to obtain SP observations at  $1.61\mu\text{m}$ . Therefore, the selection of the optimum SP channel set and interpolation scheme is based on the validation performance of Channel 2 ( $1.61\mu\text{m}$ ) rather than Channel 1 ( $0.63\mu\text{m}$ ) for SSF data. From visual inspection of regression statistics in Tables 6a and 6b, first order interpolation with SP Channel Set I (see highlighted row in Table 6a) appears to be the optimum choice for SSF validation. This choice is not anticipated since one would expect that the best agreement should be achieved for the channel set requiring the least amount of spectral extrapolation. We suspect this probably is due to insufficient match-up samples found for the SSF4 data (only 25). More match-up points are expected with the next version of the data (SSF-ED1 uses higher resolution land mask) and will be used to re-examine this conclusion. There are some water vapor absorption features at  $1.02\mu\text{m}$ , which have not been accounted for in the AERONET aerosol optical thickness. This may also affect the result. It can be examined soon since AERONET data will include wavelength at  $1.60\mu\text{m}$  and  $2.2\mu\text{m}$  in the near future (B. Holben, personal communication).

In summary, for AEROBS validation, 2<sup>nd</sup> order interpolation of the SP Channel Set I is selected, while, for SSF validation, the same SP Channel Set I but with 1<sup>st</sup> order interpolation is optimal. One additional advantage of selecting SP channel set I is that it



### 3) Sampling Approach

Sensitivity of the validation procedure to satellite data sampling approaches has also been studied. AEROS data for Channel 1 ( $0.63\mu\text{m}$ ) is used because it provides many more match-up points for validation than SSF-4 data. Four sampling approaches have been investigated, using AERONET SP Channel Set I with 2<sup>nd</sup> order interpolation as truth. The first is the  $\hat{\text{ensemble}}$  approach, which uses all AEROS aerosol optical thickness values (or 500 closest in distance, if total is more than 500) in the optimum match-up window (1 hour/100km) to determine the aerosol optical thickness mean and variance for that AERONET site and day. The second is the  $\hat{\text{best}}$  approach, where the AEROS with  $\tau_{\text{st}}$  closest to  $\tau_{\text{sp}}$  is selected from the optimum match-up window. The third is the  $\hat{\text{closest}}$  approach, where the closest (in distance) AEROS to the SP location is selected from the optimum match-up window. The last is the  $\hat{\text{ten closest}}$  approach, which is the same as the  $\hat{\text{closest}}$  but using the 10 closest (in distance) AEROS to compute mean and variance statistics (used in Ignatov et al., 1995a, b; Stowe et al., 1997). Three AERONET stations, Dry Tortugas, Bermuda, and Kaashidhoo, are used in this study.

Scatter plots and associated linear regression lines of  $\tau_{\text{st}}$  vs.  $\tau_{\text{sp}}$  for the three selected stations and four  $\tau_{\text{st}}$  sampling approaches are displayed in Fig. 3. Generally, the regression lines for the  $\hat{\text{best}}$  and  $\hat{\text{closest}}$  approaches are at the two extremes of

sufficient for a reliable statistical analysis of AEROBS data. Since AEROBS and AERONET data are not exactly collocated or invariant in space and time, it is preferable to compare  $\tau_{st}$  and  $\tau_{sp}$  averaged over some space-time window. Furthermore, the final output products of AVHRR (and TRMM/VIRS) aerosol optical thickness are in grided format (i.e., averaged over a space-time window). Thus, it is concluded that the  $\hat{\text{ensemble}}$  approach is most appropriate for validation of these aerosol products.

Although the above study is performed only for AEROBS data, it is likely that this conclusion is generally applicable to validation of other satellite sensor aerosol retrievals, such as TRMM/VIRS.

### 3. Validation Results

Using the standard procedures optimized through the above sensitivity studies, global (ensemble) validation on the 2<sup>nd</sup> generation NOAA/NESDIS operational aerosol retrieval algorithm was performed for 1998 NOAA-14/AVHRR (AEROBS) data and TRMM/VIRS (CERES-SSF4) data with AERONET observations.

The global linear regression equations predicting  $\tau_{st}$  for channels 1 (0.63 $\mu\text{m}$ ) and 2 (0.83 $\mu\text{m}$ ) of NOAA-14/AVHRR (AEROBS) from  $\tau_{sp}$  are

$$\mathbf{t}_{st}^1 = 0.062(\pm 0.015) + 0.95(\pm 0.08)\mathbf{t}_{sp}^1, \quad (2a)$$

$$\mathbf{t}_{st}^2 = 0.086(\pm 0.009) + 0.99(\pm 0.07)\mathbf{t}_{sp}^2, \quad (2b)$$

The same equations for channels 1 (0.63 $\mu$ m) and 2 (1.61 $\mu$ m) of TRMM/VIRS (CERES-SSF4) observations are

$$\mathbf{t}_{st}^1 = 0.086(\pm 0.025) + 0.84(\pm 0.14)\mathbf{t}_{sp}^1, \quad (3a)$$

$$\mathbf{t}_{st}^2 = 0.026(\pm 0.013) + 0.90(\pm 0.17)\mathbf{t}_{sp}^2, \quad (3b)$$

$$\mathbf{a}_{st} = 1.167(\pm 0.821) - 0.25(\pm 0.74)\mathbf{a}_{sp} \quad (3c)$$

with standard errors of  $\sigma_1=0.061$ ,  $\sigma_2=0.036$ , and  $\sigma_{\alpha}=1.278$  and correlation coefficients of  $R_1=0.79$ ,  $R_2=0.75$ ,  $R_{\alpha}=0.07$ . The NOAA-14/AVHRR (AEROBS) validation is much more reliable than that of TRMM/VIRS (CERES-SSF4) because there are 120 match-ups for AVHRR but only 25 for VIRS. This may be the principal reason for the difference in regression coefficients for the two satellite observations at 0.63 $\mu$ m. Also, the CERES-SSF data set is being reprocessed with improved VIRS cloud and land masks, so this validation is clearly very preliminary and is only shown for comparison to AVHRR. It will not be discussed further.

The NOAA-14/AVHRR validation at 0.63 $\mu$ m is somewhat worse than three previous validations for NOAA-9 and 11/AVHRR using ship borne sun photometers (see Ignatov et al., 1995a, b; Stowe et al., 1997). This is mainly due to differences in the validation procedures, which is a good illustration of the importance of standardizing these procedures. More subjective decisions were involved in these earlier validations. For

rejected. In the NOAA-14/AVHRR validation, only the last filter (SP records with large temporal variability) is applied to exclude rare defects in the AERONET data.

Fig. 4 shows scatter plots of the 120 AEROBS-AERONET match-up days for  $\tau_1$ ,  $\tau_2$ , and  $\alpha$  (means and standard deviations) with the corresponding linear regression equations. One can see from the figure how the  $\tau_{st}$  filtering used in the earlier validation would improve the results. The cause of these outliers (far off regression line and large  $s_{t_w}$ ) are thought to be due to cloud contamination but this has not yet been established. It is obvious from Fig. 4 (and Eqs. (2c) and (3c)) that there is larger uncertainty in the derived values of  $\alpha$  from the retrieved  $\tau$ s. Ignatov et al. (1998) and Ignatov and Stowe (this issue) have shown that random errors in  $\tau$  are amplified when deriving  $\alpha$ , particularly as  $\tau$  approaches zero. This is a result of the defining logarithm ratio

relationship ( $a = - \frac{\ln[t_1/t_2]}{\ln[I_1/I_2]}$ ). It is of interest to mention that both forward

numerical retrieval sensitivity studies (Mischenko et al., 1999) and theoretical analyses (Ignatov and Stowe, 2000) confirm that  $\alpha$  is less subject to error in the assumed aerosol retrieval model than are the  $\tau$ s from which it is derived. Thus, they conclude that  $\alpha$ , which is least sensitive to uncertainties in the atmosphere-ocean model, should be retrieved along with  $\tau$  as a second aerosol parameter. However, our validation indicates

ratio. This is clearly not substantiated by the actual observations used in our validation (bottom panel in Fig. 4). Actually, there are sufficient random errors in the retrieved  $\tau_s$  (may be from error in the aerosol retrieval model or other errors in the measurement) that are non-multiplicative and therefore do not cancel when taking their ratio (see also Higurashi et al., 2000). It appears that an algorithm, in which  $\alpha$  is derived directly from  $\tau_s$  in two separated channels, can not give quantitatively dependable values (qualitatively, perhaps). The qualitative information is still useful for separating broad categories of aerosol types (dust, haze, smoke etc.) as has been demonstrated by others (Mishchenko et al., 1999; King et al., 1999; Higurashi et al., 2000). This issue is revisited in the following section along with results of investigations to identify possible sources of systematic error in  $\tau$  as implied by the regression lines in Fig. 4. Thus, the validation procedure is shown to be a useful tool for adjusting aerosol retrieval algorithm parameters to reduce these errors (through the dash line loop in Fig.1).

#### **4. Application of the Validation Procedure to Algorithm Improvement**

##### **a) Specular Reflectance**

To eliminate the impact of specular reflectance from the rough oceanic surface, aerosol retrievals in the 2<sup>nd</sup> generation NOAA/NESDIS operational algorithm are limited to gamma ( $\gamma$ ) angles (angle between viewing angle and specular ray from the flat ocean)

up data base for both AEROBS-SP and SSF4-SP data. Accordingly, these data sets have been subjected to the validation procedure and results are summarized in Tables 7a and 7b, respectively, for AEROBS and SSF4.

The improvement of regression results for both channels of AEROBS data is minor with increases in the  $\gamma$  angle limit from  $40^\circ$  to  $60^\circ$ . However, for SSF data, the change of regression coefficients is more irregular. For both  $0.63\mu\text{m}$  and  $1.61\mu\text{m}$  channels, the  $\tau_{\text{st}}$  retrieval does appear to be sensitive to increases in the  $\gamma$  angle limit. However, all regression parameters become worse with increasing  $\gamma$  angle limit, opposite to what is expected if specular reflection were affecting the retrievals. This is again probably due to the much lower number of regression points (25 in SSF4 compared to 120 in AEROBS for  $\gamma > 40^\circ$ ) involved in the SSF4 analysis. Thus, based solely on the AEROBS analysis, it is concluded that increasing the  $\gamma$  angle limit beyond the nominal  $40^\circ$  does not significantly remove errors in aerosol optical thickness, and therefore, specular reflection of radiation beyond  $40^\circ$  is probably not a serious problem.

#### b) Calibration, Rayleigh Scattering, and Diffuse Surface Reflectance

Lowering the systematic high bias at low aerosol optical thickness in the above validation results (positive regression line intercept in Fig. 4) has been sought by checking the values of non-aerosol related elements (including calibration, Rayleigh

shown here. In Table 8, the original AEROBS operational values of calibration, Rayleigh scattering, and diffuse surface reflectance are listed as well as values after correction or adjustment based on sensitivity studies using our validation procedure. The imaginary part of the aerosol refractive index is also listed, corresponding to the discussion in the following sub-section c).

The regression statistics before and after the adjustments (or corrections) listed in Table 8 are summarized in Table 9. First, the most recent calibration slope drift correction coefficients (C.R.N. Rao, 1998 personal communication) are used to correct AEROBS reflectances. These are then used to derive new  $\tau_1$  and  $\tau_2$  values (dash line loop in Fig. 1), to which the global validation procedure is again applied. The high bias at low aerosol optical thickness (or intercept A) is lowered in both channels 1 and 2 as a result. However, the proportional error (slope B) and correlation ( $R^2$ ) are moved further away from their ideal values, while there is a small improvement in the standard error.

The Rayleigh scattering optical thickness used in our operational algorithm has been found to be in error, based on 6S radiative transfer model calculations for the exact response functions of channels 1 and 2 for NOAA-14/AVHRR (A. Ignatov, 2000 personal communication). It is too large in both channels (cf. Table 8). Using these correct values in the retrieval LUTs actually increases the bias at low aerosol optical thickness. The change of the other regression parameters (except for slope in channel 2)

Sensitivity studies have been performed with the validation procedure to identify an optimal diffuse surface reflectance that, together with the correct calibration and Rayleigh optical thickness, reduces the bias at low  $\tau$  to near zero. For each retrieval channel of AVHRR, several values of surface diffuse reflectance have been used to construct new LUTs for deriving  $\tau_1$  and  $\tau_2$  from the reflectances, to which the global validation is again applied (dash line loop in Fig.1). The values of  $\rho_{\text{dsr}}$  that yield an intercept closest to zero in each channel are shown in Table 8 (DSR heading) and the corresponding regression statistics are listed in Table 9. The new values for  $\rho_{\text{dsr}}$  for both  $0.63\mu\text{m}$  and  $0.83\mu\text{m}$  are larger than before, so less aerosol is required to match the observed reflectances. While greatly improving the bias at low  $\tau$ , it is apparent from Table 9 that the other regression statistics are only slightly degraded. The new diffuse surface reflectances are somewhat larger than expected for open oceans. This may suggest that there are coastal (e.g., shallow water) effects within 25-100km of each AERONET site or that the simple Lambertian surface model assumption, with its adjustment for diffuse glint used in the current operational algorithm (Stowe et al., 1997), has to be raised to non-physically high values in order for the retrievals to be unbiased at low aerosol optical thickness. The 6S radiation transfer model, which treats the surface reflectance as a wavy Fresnel surface with wind driven slopes, will provide our future LUTs (see Ignatov and Stowe, this issue). It thus may allow a more physically consistent value to again be used for this



### c) Imaginary Part ( $n_i$ ) of the Aerosol Refractive Index

The above improvements from non-aerosol parameters, which reduce the bias at low optical thickness, actually worsen the bias at large optical thickness (slope of regression line further away from unity). The only way to remove this bias and not affect the intercept is by adjusting parameters of the aerosol microphysical model used in the retrieval algorithm.

The imaginary part of the aerosol refractive index is the obvious first choice for adjustment since the aerosol is assumed to be non-absorbing in the current operational retrieval algorithm. If it is absorbing, as most tropospheric aerosol with large optical thickness is (e.g., Nakajima et al., 1996; Kaufman et al., 1997; Tanre et al., 1997; Nakajima and Higurashi, 1997; Mishchenko et al., 1999; Schmid et al., 1999; Dubovik and King, 2000), this would cause the sub-unity slope observed in the validation results (see Ignatov, 1995a; Stowe et al., 1997). Several values of  $n_i$  have been selected for each channel of AVHRR and new LUTs were generated. Again, the resulting aerosol optical thicknesses ( $\tau_1$  and  $\tau_2$ ) are then subjected to the global validation procedure (dash line loop in Fig. 1). The  $n_i$  that yields a slope closest to unity in each channel is 0.005 for channel 1 (0.63 $\mu\text{m}$ ) and 0.008 for channel 2 (0.83 $\mu\text{m}$ ) (cf. Table 8). A similar wavelength dependence of  $n_i$  for these two AVHRR channels has also been observed by Higurashi et al. (2000) in the validation of their two-channel AVHRR aerosol retrieval algorithm.

variability in aerosol type. The 2<sup>nd</sup> generation algorithm uses a uniform aerosol model (a log-normal distribution with a 0.1 $\mu$ m mode radius and 2.03 $\mu$ m variance and refractive index  $m=1.4-i0.0$ ) globally. The current match-up sample size for a single site of AERONET in 1998 is too small to verify this hypothesis with the validation procedure. AVHRR (AEROBS) data are being collected for 1999 and 2000 so that the regional validation can be done.

A new retrieval algorithm that accounts for regional difference in aerosol particles may be required to reduce this random error and possibly some of the systematic errors as well. As mentioned before, this kind of algorithm needs to be designed carefully to minimize the effects of  $\tau$  errors on a derived size parameter, like  $\alpha$ . There are at least two approaches. The first is to derive  $\alpha$  iteratively from a set of LUTs corresponding to different  $\alpha$  values. These LUTs should cover the range of possible atmospheric values of  $\alpha$  with sufficient resolution. The  $\alpha$  derived by using  $\tau$ s from the 2<sup>nd</sup> generation algorithm can be used as a first guess. The first guess  $\alpha$  is then used to find a closely corresponding LUT for deriving new  $\tau$ s and  $\alpha$ . This process will continue until the derived  $\alpha$  converges. The second approach is to make  $\alpha$  another dimension in a LUT and to retrieve  $\tau$ s and  $\alpha$  simultaneously. The first approach should yield more accurate  $\tau$ s by including regional size information of aerosols through  $\alpha$  on the LUTs, which in turn may reduce errors

Angstrom wavelength exponent to vary in the LUTs. This makes the retrieval algorithm sensitive to regional differences in aerosol in some senses and should reduce random error accordingly. Further discussion of this topic is outside the scope of this paper and will be presented later by its developers. The validation procedure presented here will play a critical role in assessing the performance of the new algorithm.

As a summary of this section, improvements of the validation results at low aerosol optical thickness (by adjusting non-aerosol model parameters) and at high aerosol optical thickness (by correcting the imaginary part of the aerosol refractive index) are demonstrated if one compares the scatter plots of Fig. 5 with those of Fig.4 for AEROBS data.

## **5. Some Discussions**

We feel the need for discussions on some important issues related to our validation before we draw final conclusions. First is the representativeness of the AERONET measured aerosol to the global tropospheric aerosols. Actually, very careful study on this issue has been performed recently by the AERONET scientists (e.g., Kaufman et al., 2001; Smirnov et al., this issue). They have compared AERONET measurement (spanning over 2-5 years) of  $\tau$  and size parameters (such as Angstrom wavelength  $\alpha$  and effective radius  $R_{\text{eff}}$ ) over selected AERONET marine sites in both the Pacific (Nauru,

AERONET aerosol measurements can be used to represent the climatologies of tropospheric maritime aerosols as long as the AERONET sites are carefully selected.

The next issue is how to reasonably extend the adjustment or correction we identified through sensitivity studies described in Section 4 to global implementation, especially for the diffuse surface reflectance and the aerosol model used in the retrieval algorithm. Regional validation based on sufficient match-up samples is a feasible next step. Three-years of match-up data (1998, 1999, and 2000) are being collected now for reliable regional validation. First, the diffuse surface reflectance determined from the sensitivity studies for the AERONET sites that are surrounded by deep water and less waves will be analyzed to determine an appropriate value of the adjustment that can be applied globally in the algorithm. For example, Lanai is a good candidate site due to the fact that it is surrounded by deep water, is sheltered from wind by the surrounding islands, receives very little rain, and is not known for large waves (Price, 1983; Smirnov et al., this issue). Second, baseline AERONET sites can be further divided into different categories according to the prevailing origins of the aerosols over them (such as maritime, mineral dust, biomass burning, and urban/industrial, or the mix of them) and the aerosol properties derived from the skylight analyses. Sensitivity studies used in the global validation for aerosol model adjustment can be similarly applied to each category of sites. This will allow us to evaluate the aerosol regional representativeness of a retrieval

for a retrieval algorithm that is sensitive to some kind of aerosol type (such as our 3<sup>rd</sup> generation algorithm). The purpose of the regional validation is to identify and document this variable performance of a retrieval algorithm in a quantitative way. The validation procedures proposed in this paper form a basis for achieving this purpose.

One may have noticed that we have basically made two kinds of manipulation to maximize the agreement between the satellite retrieval and the ground truth. The first is the optimization procedures based purely on the statistic regression analyses. The second is the fine adjustments to the intercept and slope of the linear regression formula based on solid physical rationale, which is illustrated by using the following linearized single scattering approximation of the radiative transfer equation (see also Stowe et al., 1997):

$$\tau_{ST} = 4m_s m_h \frac{\rho - \rho^R - \rho^S T}{\omega P^A}, \quad (4)$$

where  $\rho$  is an apparent reflectance of the ocean-atmosphere system;  $\rho^R$  is the Rayleigh scattering contribution;  $\rho^S$  is the diffuse surface reflectance;  $T$  is the total atmospheric transmittance;  $P^A$  and  $\omega$  are the aerosol phase function and single scattering albedo;  $\mu_s$  and  $\mu_v$  are the cosine of solar and view zenith angles. The errors in aerosol optical thickness  $\tau_{ST}$  may result from  $\rho^S$ ,  $P^A$ , and  $\omega$  since all other terms are reasonably well known. Note that  $\rho$  and  $\rho^S$  participate in the equation as additive terms, and  $(\omega P^A)$  as multiplicative one. This suggests that the non-zero intercept ( $A \neq 0$ ) in the regression

validation approach is based on a combination of purely statistical regression with fine adjustments to the intercept and slope of the regression formula based on physical rationale.

One can see that after sufficient correlation has been established from the statistical analysis, further delicate adjustment on the slope and intercept helps us identify possible sources of error and reduce them in the retrieval algorithm with only minor degradation on the correlation as shown in Table 9. We think this combination of pure statistical analysis with the intercept and slope adjustments (supported by physical rationale) is the unique aspect of the validation procedure used in the paper. Of course, we have to admit that more investigations are needed to justify the quantity of the adjustment suggested from the sensitivity studies. This is very difficult to do using the Dave radiative transfer model used in our 2<sup>nd</sup> generation operational algorithm due to its limitations. This is better done with our 3<sup>rd</sup> generation algorithm based on the more complete 6S radiative transfer model, which will give more precise answers to these kinds of questions. The new features in the 6S code, such as a wavy Fresnel surface with wind driven slopes and more widely used aerosol model (e.g., bi-model log-normal distribution), will make the further physical quantitative studies possible. It can be seen that the validation procedures adopted in this paper and the data sets collected form a basis for testing the new algorithm.

been developed. Based on detailed sensitivity studies, a 1 hour/100km time/space match-up window has been identified to assure sufficient collocation and comparable space-time variance statistics for aerosols measured from two different platforms (satellite and ground). Also, the ensemble method of averaging on this time/space scale was found to be best for identifying biases in satellite aerosol retrieval products (such as optical thickness and Angstrom wavelength exponent) especially important for climate research. This space/time scale is also close to the scales used when gridding these pixel level retrieval products for statistical analyses. For validation of aerosol retrievals from NOAA-14/AVHRR with AERONET measurements, the final result is not sensitive to the choice of interpolation scheme and SP channel sets. Thus, for historical continuation, a second order polynomial fit in logarithm of wavelength to the four SP channels is used to determine the AERONET optical thickness for the two AVHRR channels. For validation of TRMM/VIRS (CERES-SSF4) retrievals, the same four SP channels, but with first order polynomial interpolation, are optimal, even though extrapolation is required for the 1.61 $\mu$ m channel.

This global validation package has been applied to the 2<sup>nd</sup> generation NOAA/NESDIS operational aerosol retrieval algorithm by using 1998 observations from NOAA-14/AVHRR (AEROBS) and TRMM/VIRS (CERES-SSF4). It has been shown to be useful for improving the satellite aerosol retrieval by making adjustments to aerosol

(0.83 $\mu\text{m}$ ) are summarized in Table 10. A similar summary for 1998 VIRS (CERES-SSF4)  $\tau$  data for channel 1 (0.63 $\mu\text{m}$ ) and 2 (1.61 $\mu\text{m}$ ) is given in Table 11. Values, both before and after correcting and adjusting non-aerosol model parameters and the imaginary part of the aerosol refractive index, are listed for comparison.

The NOAA-14/AVHRR (AEROBS)  $\tau$  values for mean conditions are biased high by 0.05 and 0.08, with random errors of 0.08 and 0.05, at 0.63 $\mu\text{m}$  and 0.83 $\mu\text{m}$ , respectively. Correspondingly, the TRMM/VIRS (CERES-SSF4)  $\tau$  values for mean conditions are biased high by 0.06 and 0.02, with random errors of 0.06 and 0.04, at 0.63 $\mu\text{m}$  and 1.61 $\mu\text{m}$ , respectively. After corrections and adjustments, the biases in both channels of AVHRR and VIRS have been reduced significantly and are close to zero, although random error is almost unchanged. The effective Angstrom wavelength exponent  $\alpha$ , derived directly from the  $\tau$ s, has been shown to be poorly correlated both before and after adjustments, indicating that random error in the  $\tau$  measurement (possibly related to aerosol model parameter variations or cloud/surface reflectance contamination) needs to be reduced. The results of TRMM/VIRS validation need to be viewed with caution due to the small match-up sample size resulting from use of a preliminary version 4 of the CERES-SSF data set. However, it is interesting to see that the algorithm adjustments and statistical results of SSF4 aerosol validation are not in conflict with those from AVHRR/AEROBS even though the sample size of SSF4 match-ups is much smaller



analysis, which will help to identify algorithm biases due to regionally different aerosol types. These global validation studies suggest that the random errors in the current NOAA/NESDIS operational aerosol retrieval algorithm could very well be due to regional differences in aerosol particles. These differences are to be included in the 3<sup>rd</sup> generation algorithm, currently under development at NESDIS. The new algorithm, similar to those already developed by Mishchenko et al. (1999), Higurashi and Nakajima (1999), and Tanre et al. (1997), will derive aerosol optical thickness and a size parameter (such as Angstrom wavelength exponent or effective radius) from two (or three for NOAA-16) channels, used dependently, rather than independently. This makes the retrieval sensitive to regional differences in aerosol, particularly with respect to particle size. The validation procedure presented here will be applied to the results from the 3<sup>rd</sup> generation algorithm to evaluate and document its performance. We also hope the standardized validation procedure presented here can be used by other research groups for their algorithm validation, which will make algorithm inter-comparison much more meaningful in the future. The archived match-up data set is available for algorithm inter-comparison upon request from the authors.

### **Acknowledgments**

We express our deep gratitude to the AERONET project leader Dr. Brent Holben at

TRMM/VIRS (CERES-SSF4) 1998 data. Discussions with Dr. Alexander Ignatov at NOAA/NESDIS also were of great benefit. Two anonymous reviewers' comments were very constructive for improvements to the manuscript. This work was funded by NASA Langley through contract #L90987C managed by Drs. Bruce Wielicki and Bruce Barkstrom, CERES science team managers, and by NOAA/NESDIS through a grant from the ocean remote sensing program managed by Dr. Lee Dantzer.

### References

- Beyer, H. W., Editor, 1991: *CRC Standard Mathematical Tables and Formulae*, 29<sup>th</sup> Edition, Probability and Statistics (chapter 8), CRC Press, Inc., 609pp.
- Charlson, R. J., S. E. Schwartz, J. M. Hales, R. D. Cess, J. A. Coakley Jr., J. E. Hansen, and D. J. Hofmann, 1992: Climate forcing by anthropogenic aerosols, *Science*, **255**, 423-430.
- Chu, D. A., Y. J. Kaufman, L. A. Remer, and B. N. Holben, 1998: Remote sensing of smoke from MODIS airborne simulator during the SCAR\_B experiment, *J. Geophys. Res.*, **103**, 31,979-31,987.
- Chu, D. A., Y. J. Kaufman, D. Tanre, and B. N. Holben, 1999: Smoke optical properties derived from airborne measurements in Amazonia, paper presented at *ALPS'99* Cent,

Aerosol Robotic Network (AERONET) sun and sky radiance measurements, *J. Geophys. Res.*, **105**, 9791-9806.

Dubovik, O., and M. King, 2000: A flexible inversion algorithm for retrieval of aerosol optical properties from Sun and sky radiance measurements, *J. Geophys. Res.*, **105**, 20,673-20,696.

Durkee, P. A., K. E. Nielsen, P. J. Smith, P. B. Russell, B. Schmid, J. M. Livingston, B. N. Holben, C. Tomasi, V. Vitale, D. Collins, R. C. Flagan, J. H. Seinfeld, K. J. Noone, E. Ostrom, S. Gasso, D. Hegg, L. M. Russell, T. S. Bates, and P. K. Quinn, 1999: Regional aerosol optical depth characteristics from satellite observations: ACE-1, TARFOX and ACE-2 results, *Tellus*, in press.

Goloub, P. D., D. Tanre, J. L. Deuze, M. Herman, A. Marchand, and F. M. Breon, 1999: Validation of the first algorithm applied for deriving the aerosol properties over the ocean using POLDER/ADEOS measurements, *IEEE Trans Geosci. Remote Sens.*, **37**, 1586-1596.

Hansen, J. E., and A. A. Lacis, 1990: Sun and dust versus greenhouse gases: An assessment of their relative roles in global climate change, *Nature*, **346**, 713-719.

Hansen, J. E., M. Sato, R. Ruedy, A. Lacis, and V. Oinas, 2000: Global warming in the 21<sup>st</sup> century: an alternate scenario, *Proc. Nat. Acad. Sci.*, **97**, 9875-9880.

Higurashi, A., T. Nakjima, B. N. Holben, A. Smirnov, R. Frouin, and B. Chatenet, 2000:

- Holben, B. N., T. F. Eck, I. Slutsker, D. Tanre, J. P. Buis, A. Setzer, E. Vermote, J. A. Reagan, Y. J. Kaufman, T. Nakajima, F. Lavenu, I. Jankowiak, and A. Smirnov, 1998: AERONET-A federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, **66**, 1-16.
- Husar, B. R., J. M. Prospero, and L. L. Stowe, 1997: Characterization of tropospheric aerosols over the oceans with the NOAA advanced very high resolution radiometer optical thickness operational product, *J. Geophys. Res.*, **102**, 16,889-16,909.
- Ignatov, A. M., L. L. Stowe, S. Sakerin, G. Korotaev, 1995a: Validation of the NOAA/NESDIS satellite aerosol product over the North Atlantic in 1989, *J. Geophys. Res.*, **100**, 5123-5132.
- Ignatov, A. M., L. L. Stowe, R. Singh, D. Kabanov, and I. Dergileva, 1995b: Validation of NOAA/AVHRR aerosol retrievals using sun-photometer measurements from RV Akademik Vernadsky in 1991, *Adv. Space Res.*, **16**(10), 95-98.
- Ignatov, A. M., L. L. Stowe, and R. Singh, 1998: Sensitivity study of the Angstrom exponent derived from AVHRR over the oceans, *Adv. Space Res.*, **21**, 439-442.
- Ignatov, A. M., and L. L. Stowe, 2000: Physical basis, premises, and self-consistency checks of aerosol retrievals from TRMM/VIRS, *J. Appl. Meteor.*, **39**, 2259-2277.
- IPCC, 1996: *Climate Change 1995. The Science of Climate Change*, Cambridge University Press, 572pp.

tropospheric aerosol and atmospheric correction for the aerosol effect, *J. Geophys. Res.*, **102**, 16,815-16,830.

Kaufman, Y. J., A. Smirnov, B. N. Holben, and O. Dubovik, 2001: Baseline maritime aerosol: methodology to derive the optical thickness and scattering properties, *Geophys. Res. Lett.* (submitted).

King, M. D., Y. J. Kaufman, D. Tanre, and T. Nakajima, 1999: Remote sensing of tropospheric aerosols from space: past, present, and future, *Bull. Am. Meteorol. Soc.*, **80**, 2229-2259.

Lacis, A. A., and M. I. Mishchenko, 1995: Climate forcing, climate sensitivity, and climate response: A radiative modeling perspective on atmospheric aerosols, in *Aerosol Forcing of Climate*, R. Charlson and J. Heintzenberg, eds., pp.11-42, Wiley, New York.

Mishchenko, M. I., I. V. Geogdzhayev, B. Cairns, W. B. Rossow, and A. Lacis, 1999: Aerosol retrievals over the oceans by use of channels 1 and 2 AVHRR data: Sensitivity analysis and preliminary results, *Appl. Opt.*, **38**, 7325-7341.

Nakajima, T., G. Tonna, R. Rao, P. Boi, Y. Kaufman, and B. Holben, 1996: Use of sky brightness measurements from ground for remote sensing of particulate polydispersions, *Appl. Opt.*, **35**, 2672-2686.

Nakajima T., and A. Higurashi, 1997: AVHRR remote sensing of aerosol optical

- Price, S., 1983: *Atlas of Hawaii*, University of Hawaii Press.
- Schmid, B., J. Michalsky, R. Halthore, M. Beauharnois, L. Harrison, J. Livingston, P. Russell, B. Holben, T. Eck, and A. Smirnov, 1999: Comparison of aerosol optical depth from solar radiometers during fall 1997 ARM intensive observation period, *Geophys. Res. Lett.*, **26**, 2725-2728.
- Smirnov, A., B. N. Holben, T. F. Eck, O. Dubovik, and I. Slutsker, 2000: Cloud screening and quality control algorithms for the AERONET data base, *Remote Sens. Environ.*, **73**, 337-349.
- Smirnov, A., B. N. Holben, Y. J. Kaufman, O. Dubovik, T. F. Eck, I. Slutsker, C. Pietras, and R. Halthore, 2001: Optical properties of atmospheric aerosol in maritime environments, *J. Atmos. Sci.* (this issue).
- Stowe, L. L., A. M. Ignatov, and R. R. Singh, 1997: Development, validation, and potential enhancements to the second-generation operational aerosol product at the National Environmental Satellite, Data, and Information Service of the National Oceanic and Atmospheric Administration, *J. Geophys. Res.*, **102**, 16,923-16,934.
- Stowe, L. L., and H. Jacobowitz, 1997: The AVHRR Pathfinder Atmosphere (PATMOS) Project, *Proc. 9<sup>th</sup> Atmospheric Radiation Symposium*, pp.196-197, AMS, Boston, MA.
- Tanre, D., M. Herman, and Y. J. Kaufman, 1996: Information on the aerosol size

Tanre, D., Y. J. Kaufman, M. Herman, and S. Matoo, 1997: Remote sensing of aerosol properties over oceans using the MODIS/EOS spectral radiances. *J. Geophys. Res.*, **102**, 16,971-16,988.

### Figure Captions:

Fig. 1. Flow chart of global aerosol validation procedure.

Fig. 2. Plots of regression coefficients (intercept, slope, standard error, and square of correlation coefficient) for 20 different match-up window sizes at the Dry Tortugas AERONET site for AVHRR (AEROBS) aerosol optical thickness from the  $0.63\mu\text{m}$  channel.

Fig. 3. Scatter plots and regression lines for  $\tau$  at  $0.63\mu\text{m}$  for four averaging approaches (see text) applied to NOAA-14/AVHRR (AEROBS) 1998 operational data with a match-up window of 1 hour/100km at Dry Tortugas, Bermuda, and Kaashidhoo AERONET sites. Regression formula  $y = Ax + B$  and correlation parameter,  $R^2$ , have also been presented for the four approaches, where  $x$  represents  $\tau_{\text{sp}}$  and  $y$  represents  $\tau_{\text{st}}$ . Colors are used to identify the match-up points, regression lines, and regression formulas from the four approaches.

Fig. 4. Scatter plots of  $\tau_1$ ,  $\tau_2$ , and  $\alpha$  and linear regression lines from the optimum global (all 7 AERONET sites from Table 1) validation of 1998 AVHRR (AEROBS) data which used the operational (1996) calibration scheme. Horizontal and vertical error bars are +/- one standard deviation long.

Fig. 5. Same as Fig. 4 but with an improved AVHRR calibration scheme (1998), and a



Table 1. Selected thirteen AERONET island stations and their location (latitude and longitude) for our global aerosol retrieval validation. The highlighted stations are those picked with sufficient Level 2 data in 1998 for our validation.

No	Stations	Latitude, Longitude
<b>1</b>	<b>Andros Island</b>	<b>24.68, -77.78</b>
2	Ascension Island	-7.97, -14.40
<b>3</b>	<b>Bahrain</b>	<b>26.32, 50.50</b>
4	Barbados	13.17, -59.50
<b>5</b>	<b>Bermuda</b>	<b>32.37, -64.68</b>
6	Cape Verde	16.72, -22.93
7	Dakar	14.38, -16.95
<b>8</b>	<b>Dry Tortuga</b>	<b>24.60, -82.78</b>
9	Guadeloup	16.32, -61.50
<b>10</b>	<b>Kaashidhoo</b>	<b>4.95, 73.45</b>
<b>11</b>	<b>Lanai</b>	<b>20.82, 156.98</b>
12	St. Nicolas	33.25, -119.49
<b>13</b>	<b>Surinam</b>	<b>5.78, -55.20</b>

Table 2. Selected time/space match-up windows used in sensitivity studies for determining optimal match-up window.

Time Window (hour)	Space Window (Radius of the Match-up Circle Around an AERONET Station)				
	100 (km)	200 (km)	300 (km)	400 (km)	500 (km)
+/- 1	W11	W12	W13	W14	W15
+/- 2	W21	W22	W23	W24	W25
+/- 3	W31	W32	W33	W34	W35
+/- 4	W41	W42	W43	W44	W45



Table 5a. Regression statistics for validation of AEROBS data using two SP channel sets interpolated with a 1<sup>st</sup> order polynomial in natural logarithm of wavelength.

Satellite Wavelength	SP Channel Set	A Value	$\Delta A$ Value	B Value	$\Delta B$ Value	$\sigma$ Value	$R^2$ Value
<b>0.63</b>	<b>I</b>	<b>0.0591</b>	<b>0.0147</b>	<b>1.0113</b>	<b>0.0870</b>	<b>7.7959E-2</b>	<b>0.5315</b>
( $\mu\text{m}$ )	II	0.0583	0.0148	1.0166	0.0876	7.8002E-2	0.5310
0.83	I	0.0862	0.0090	0.9831	0.0721	5.1038E-2	0.6099
( $\mu\text{m}$ )	II	0.0855	0.0089	0.9902	0.0711	5.0383E-2	0.6199

Table 5b. Same as above but for 2<sup>nd</sup> order polynomial interpolation.

Satellite Wavelength	SP Channel Set	A Value	$\Delta A$ Value	B Value	$\Delta B$ Value	$\sigma$ Value	$R^2$ Value
<b>0.63</b>	<b>I</b>	<b>0.0623</b>	<b>0.0148</b>	<b>0.9514</b>	<b>0.0841</b>	<b>7.9069E-2</b>	<b>0.5181</b>
( $\mu\text{m}$ )	II	0.0627	0.0146	0.9682	0.0842	7.8388E-2	0.5263
0.83	I	0.0862	0.0090	0.9913	0.0726	5.1016E-2	0.6102
( $\mu\text{m}$ )	II	0.0862	0.0090	0.9907	0.0726	5.1036E-2	0.6099

Table 6a. Same as Table 5a but for TRMM/VIRS (CERES-SSF4) data.

Satellite Wavelength	SP Channel Set	A Value	$\Delta A$ Value	B Value	$\Delta B$ Value	$\sigma$ Value	$R^2$ Value
0.63 ( $\mu\text{m}$ )	I	0.0858	0.0253	0.8359	0.1361	6.0892E-2	0.6213
	II	0.0890	0.0251	0.8134	0.1337	6.1265E-2	0.6166
	III	0.0890	0.0251	0.8112	0.1336	6.1320E-2	0.6159
1.61 ( $\mu\text{m}$ )	<b>I</b>	<b>0.0255</b>	<b>0.0131</b>	<b>0.9002</b>	<b>0.1679</b>	<b>3.6199E-2</b>	<b>0.5556</b>
	II	0.0314	0.0133	0.7797	0.1608	3.8186E-2	0.5055
	III	0.0312	0.0132	0.7964	0.1634	3.8084E-2	0.5081

Note: Match-up data at Dry Tortugas has been eliminated in this regression computation since there is no SP observation on 1.61  $\mu\text{m}$  channel at the station.

Table 6b. Same as Table 5b but for TRMM/VIRS (CERES-SSF4) data.

Satellite Wavelength	SP Channel Set	A Value	$\Delta A$ Value	B Value	$\Delta B$ Value	$\sigma$ Value	$R^2$ Value
0.63 ( $\mu\text{m}$ )	I	0.0910	0.0257	0.7917	0.1354	6.2746E-2	0.5978
	II	0.0841	0.0258	0.8425	0.1388	6.1339E-2	0.6157
	III	0.0862	0.0255	0.8335	0.1370	6.1254E-2	0.6167
1.61 ( $\mu\text{m}$ )	I	0.0452	0.0115	0.5520	0.1188	3.8994E-2	0.4844
	II	0.0378	0.0124	0.6477	0.1360	3.8536E-2	0.4964

Table 7a. Regression statistics for determining the sensitivity of AEROBS retrieval errors to specular reflectance.

Satellite Channels	$\gamma$ angle (degree)	A Value	B Value	$\sigma$ Value	$R^2$ Value
	> 40	0.0623	0.9514	7.9069E-2	0.5181
0.63 $\mu\text{m}$	> 50	0.0607	0.9661	8.0571E-2	0.5377
	> 60	0.0612	0.9680	8.0389E-2	0.5459
	> 40	0.0862	0.9913	5.1016E-2	0.6102
0.83 $\mu\text{m}$	> 50	0.0860	1.0023	5.0504E-2	0.6407
	> 60	0.0844	1.0279	5.1070E-2	0.6528

Table 7b. Same as 7a but for SSF4 data.

Satellite Channels	$\gamma$ angle (degree)	A Value	B Value	$\sigma$ Value	$R^2$ Value
	> 40	0.0858	0.8359	6.0892E-2	0.6213
0.63 $\mu\text{m}$	> 50	0.0985	0.6690	5.2072E-2	0.6229
	> 60	0.0982	0.6621	5.7342E-2	0.5976
	> 40	0.0255	0.9002	3.6199E-2	0.5556
1.61 $\mu\text{m}$	> 50	0.0363	0.5753	2.7938E-2	0.5017

Table 8. Corrections and adjustments to the AVHRR calibration and operational algorithm to reduce positive bias at low  $\tau$  (intercept  $> 0$ ) and negative bias at large  $\tau$  (slope  $< 1$ ).

CALIBRATION SLOPE (CS)		
(d = day after launch)		
Channel	Operation (1996)	Correction (1998)
0.63 $\mu\text{m}$	$S1=0.109+2.32 \times 10^{-5}d$	$S1=0.1107+1.35 \times 10^{-5}d$
0.83 $\mu\text{m}$	$S2=0.129+3.73 \times 10^{-5}d$	$S2=0.1343+1.33 \times 10^{-5}d$
RAYLEIGH OPTICAL THICKNESS (ROT)		
Channel	Operation	Correction
0.63 $\mu\text{m}$	0.0607	0.0554
0.83 $\mu\text{m}$	0.0205	0.0180
DIFFUSE SURFACE REFLECTANCE (DSR)		
Channel	Operation	Adjustment
0.63 $\mu\text{m}$	0.002	0.01
0.83 mm	0.0005	0.0006
IMAGINARY PART OF THE AEROSOL REFRACTIVE INDEX (IPARI)		

Table 9. Regression statistics for AEROBS data before and after sequentially adjusting calibration and aerosol retrieval algorithm parameters (see Table 8) to reduce retrieval errors.

AVHRR Channels	adjusted parameter	A Value	$\Delta A$ Value	B Value	$\Delta B$ Value	$\sigma$ Value	$R^2$ Value
0.63 ( $\mu\text{m}$ )	operation	0.0623	0.0148	0.9512	0.0841	0.07907	0.5180
	1998 CS	0.0332	0.0145	0.8404	0.0821	0.07716	0.4683
	ROT	0.0654	0.0144	0.8465	0.0819	0.07707	0.4725
	DSR	0.0030	0.0147	0.8308	0.0838	0.07875	0.4525
	IPARI	-0.0019	0.0174	0.9903	0.0987	0.09284	0.4579
0.83 ( $\mu\text{m}$ )	operation	0.0861	0.0090	0.9913	0.0726	0.05101	0.6103
	1998 CS	0.0538	0.0079	0.8578	0.0641	0.04506	0.6004
	ROT	0.0679	0.0076	0.8219	0.0616	0.04326	0.5995
	DSR	0.0173	0.0083	0.8199	0.0666	0.04679	0.5601
	IPARI	0.0178	0.0099	0.9996	0.0803	0.05639	0.5658



Table 10. Systematic and random errors for 1998 AEROBS  $\tau$  data before and after making calibration, Rayleigh optical thickness, surface diffuse reflectance, and imaginary part of the aerosol refractive index corrections and adjustments listed in Table 8.

|-----Systematic Errors-----|

Correction Status	Channel ( $\mu\text{m}$ )	Minimum ( $\tau=0.00$ )	Mean ( $\tau=0.15@ \lambda_1$ ) ( $\tau=0.11@ \lambda_2$ )	Maximum ( $\tau=1.00$ )	Random Error (+/-)
Before Correction	$\lambda_1=0.63$	+0.06	+0.05	+0.01	0.08
	$\lambda_2=0.83$	+0.09	+0.08	+0.08	0.05
After Correction	$\lambda_1=0.63$	-0.01	-0.003	-0.01	0.08
	$\lambda_2=0.83$	+0.02	+0.02	+0.02	0.05

Table 11. Same as Fig.10 but for 1998 TRMM/VIRS (CERES-SSF4)  $\tau$  data before and after making Rayleigh optical thickness, surface diffuse reflectance, and imaginary part of the aerosol refractive index corrections and adjustments (no corrections were made to VIRS calibration and to Rayleigh optical thickness for channel 1.61 $\mu$ m). The errors after adjustments in 0.63 $\mu$ m channel are not optimized since adjustments were taken directly from the AEROBS analyses. Surface diffuse reflectance before and after adjustment for channel 1.61 $\mu$ m is 0.000 and 0.002, respectively, while the imaginary part of the aerosol refractive index is adjusted from 0.000 to 0.015.

|-----Systematic Errors-----|

Correction Status	Channel ( $\mu$ m)	Minimum ( $\tau=0.00$ )	Mean ( $\tau=0.16@ \lambda_1$ ) ( $\tau=0.07@ \lambda_2$ )	Maximum ( $\tau=1.00$ )	Random Error (+/-)
Before Correction	$\lambda_1=0.63$	+0.09	+0.06	-0.07	0.06
	$\lambda_2=1.61$	+0.03	+0.02	-0.07	0.04
After Correction	$\lambda_1=0.63$	+0.03	+0.04	+0.08	0.08
	$\lambda_2=1.61$	+0.004	+0.005	+0.01	0.04

# Flow Chart for Aerosol Validation






Fig. 1


**A-Plot (Intercept)**




**B-Plot (Slope)**






**Sigma-Plot**






**R\_Square-Plot**






**Fig. 2**



**Fig. 3**



**Fig. 4**



**Fig. 5**